



# Percutaneous Pain Procedures in Patients Who Underwent Lumbar Disc Herniation Surgery: Is It an Important Tool in the Management of Post-Surgical Ongoing Pain?

Furkan DIREN<sup>1</sup>, Omer OZDEMIR<sup>1</sup>, Osman BOYALI<sup>1</sup>, Yeliz BAHAR-OZDEMIR<sup>2</sup>, Serdar KABATAS<sup>1</sup>, Erdinc CIVELEK<sup>1</sup>

<sup>1</sup>University of Health Sciences, Gaziosmanpasa Training and Research Hospital, Department of Neurosurgery, Istanbul, Turkey

<sup>2</sup>University of Health Sciences, Sultan Abdulhamid Han Training and Research Hospital, Department of Physical Medicine and Rehabilitation, Istanbul, Turkey

**Corresponding author:** Omer OZDEMIR ✉ dromerozdemir1982@gmail.com

## ABSTRACT

**AIM:** To evaluate the efficacy of percutaneous pain interventions in patients who previously underwent lumbar disc herniation surgery.

**MATERIAL and METHODS:** We retrospectively analyzed 48 patients with persistent / recurring complaints who underwent lumbar disc surgery (LDS) and were treated with percutaneous interventions. They were grouped into recurrent disc herniations (RDHs) and other disc/vertebral pathologies (ODVP). Moreover, patients were evaluated as those who received transforaminal injection (TFI) with facet blockage (FB) and who received both caudal injection (CI) and TFI in addition to FB. Patients were evaluated using Oswestry Disability Index (ODI) and visual analog scale (VAS).

**RESULTS:** Between the recurrent and ODVP groups, preoperative, at 1-hour postoperative, and at 6-month postoperative ODI ( $p=0.867$ ,  $p=0.055$ ,  $p=0.892$ ) and VAS ( $p=0.902$ ,  $p=0.136$ ,  $p=0.462$ ) scores did not show a statistically significant difference, respectively. Additionally, in the comparison of patients who underwent FB+TFI+CI and only FB+TFI, there was no statistically significant correlation between preoperative and 6-month postoperative ODI ( $p = 0.284$ ) and VAS ( $p=0.248$ ) scores in both recurrent and ODVP groups, respectively. The success rates at the 3rd and 6th months of patients with RDH and ODVP were 47.61% (10/21) and 42.85% (9/21) and 70.37% (19/27) and 63.96% (17/27), respectively.

**CONCLUSION:** There was no statistically significant difference in ODI and VAS scores between recurrent and ODVP groups. The clinical success rate was numerically better in the ODVP group. Thus, we suggest that co-administration of TFI and CI did not significantly contribute to our clinical outcome.

**KEYWORDS:** Back pain, Management, Postsurgical

## INTRODUCTION

After lumbar discectomy (LD), low back pain (LBP) and radicular pain can recur in some patients, and the pain does not completely improve. Therefore, persistent pain after spine surgery is a common condition, and the prevalence rates range between 10% and 40% (13). While some patients with pain require re-operation, non-surgical treatments are

beneficial for some. Recurrent lumbar disc herniations (RLDHs) are the most common condition requiring re-operation (5,27). RLDHs are considered as herniations at the same level and from the same side within the first 6 mo after LD and have an incidence rate of 2%–25% (5,27).

Other conditions requiring re-operation to re-evaluate the patients with persistent/ recurring complaints after LD are

epidural fibrosis, adhesive arachnoiditis, isolated lateral spinal stenosis, and iatrogenic segmental instability (3,9,24,36). Among these pathologies, epidural fibrosis and segmental instability are the most common (3,9,24,36). The incidence of epidural fibrosis after LD varies between 18% and 37%, depending on the surgical technique (2). Different complications and outcomes have been reported in various studies (8,22,25). Although LD is the most common treatment for RLDHs requiring re-operation, anterior/ posterior interbody fusion and endoscopic minimally invasive interventions are also applied. Since the complication rates are higher and pain control is lower in patients who undergo re-operation, patient satisfaction is lower than in the first operation (3,27). Regarding pain is predominant without new neurological deficits in most patients whose complaints recur due to RLDH/ other reasons and require treatment (19,31). Therefore, pain management gains importance in the treatment of such patients. In pain management, epidural injections (EIs) with much lower complication rates and patient costs are performed (19,31). EIs are especially important in providing pain control in patients who are obese, have comorbidities, and are predicted to be at high risk for secondary surgical intervention (19,31). EIs can be administered through three different access routes: transforaminal, interlaminar, and caudal (19,31). With transforaminal injectiin (TFI), steroids and local anesthetic agents are administered to the proximal part of the dorsal root ganglion and nerve root (19,31). Local anesthetic agents provide pain control by regulating nerve conduction with “neuroversion” in the epidural area (4). The epidural drug mixture is administered to the anterior epidural area by spreading from the distal to proximal by caudal injection (CI) (19). It is thought that steroids provide anti-inflammatory effects by inhibiting inflammatory mediators around the epidural area and neural tissue (19,31).

This study aimed to evaluate the efficacy of TFI with facet denervation (radiofrequency median nerve block) or transforaminal and caudal epidural injection applications with facet denervation of patients who were divided into two groups [recurrence and other discovertebral pathologies (ODVP)] in lumbar magnetic resonance images (MRIs) of patients with evolving low back and leg pain complaints after LD.

## ■ MATERIAL and METHODS

Between January 2016 and January 2020, patients who were referred to the neurosurgery clinic with complaints of low back and leg pain and had a history of LDS and whose complaints were found to be related to the operative level included in the study group of interventional pain treatment procedures.

A list of patients who underwent interventional pain treatment was obtained using the hospital database. It was shown that patients' complaints were related to the level at which they were operated for lumbar disc herniation (LDH) by neurological examination and neuroradiological findings. In addition to the patient information system of the hospital, the data bank of the national health system was also used in this retrospective study. After the patient data were collected, 48 patients who could be reached by phone and were in accordance with the criteria were included in the study.

Institutional ethics approval was obtained from the local ethics committee (approval date: 26.05.2021, project number: 283). Oral and written informed consent was obtained from all patients. The study was conducted in accordance with the principles of the Declaration of Helsinki.

Interventional pain management was applied to patients who had low back and leg pain that was similar to that before LD, had neurological examination findings related to the operative level, and had hemipartial or total laminectomy defect in radiologic images that were related to the neurological findings and did not benefit from at least 4 weeks of medical treatment and/or physical therapy. The accepted neurologic examination findings are positivity of neurological provocation tests, hypoesthesia in the appropriate dermatome area,  $\leq 20\%$  strength loss, and limitation of lower back movements. Since TFI is less effective in patients with advanced neural compression, such as extruded disc herniation, it was not applied to patients with advanced neural compression and  $>20\%$  newly developed muscle strength loss (10,15,20). Radiological findings at the level of laminectomy defect in lumbar MRI showed RLDH, epidural fibrosis, osteophyte formation of the vertebral corpus, facet joint degeneration and micro-instability. Patients who could not undergo clinical follow-up, patients with very large disc herniation, a central canal diameter  $<6$  mm, spondylolisthesis more than grade I, severe instability on dynamic lumbar lateral radiographs, and history of vascular stent within 6 months; patients who could not complete the procedure because of respiratory distress in the prone position (patients with lung problems, such as chronic obstructive pulmonary disease); those who were known to have a hypersensitivity reaction to the local anesthetic and steroid; and pregnant patients were excluded from the study.

The patients were grouped into RLDs and ODVP (epidural fibrosis, micro-instability, lateral recess stenosis, central canal stenosis) according to MRIs. Radiofrequency thermo-ablation (RFT) was performed on the facet joints of all patients. Patients in both groups were divided into two subgroups as those who received TFI and those who received TFI + CI. Routinely performing facet denervation in patients aims to control LBP originating from the facet joint, apart from LDHs. This also contributed to the standardization of patients in the study. CI was performed in patients with multiple LDHs without radicular pressure on MRI, except at the lumbar level where they had undergone lumbar disc herniation surgery (LDHS). Combined treatment was performed to apply the treatment that would relieve the patients' low back and leg pain in the best way possible. Pain status was assessed using the Oswestry Disability Index (ODI) and leg visual analog scale (VAS) before and 1 h after the procedure. Six months after the procedure, patients were called by telephone, and their final status was evaluated with the ODI and leg VAS. Patients whose complaints improved after pain treatment procedures and who did not require surgical treatment during the follow-up were called again after 14–16 months, and the state of well-being was determined using ODI and VAS. Patients with  $<50\%$  pain complaints at the 6th month and 1st year postoperatively were recorded. For phone interviews, a patient

list including patient name-surname, identity number of the Republic of Turkey, phone number, date of surgery, date of interventional pain treatment, and preintervention complaints was created. Phone calls were made by the newly recruited resident physician who did not know the patients and did not have detailed information about the procedures.

Informed consent was obtained from the patient and his relatives about the procedure to be applied and the use of data before the interventional pain treatment.

A single-dose 1 g cefazolin was administered to the patients 30 min before the procedure. The patients were placed on the operating table in a neutral position prone on a silicone gel roll. A local anesthetic injection containing 80 mL of lidocaine, 0.05 mg epinephrine, and 50 mg of bupivacaine was administered under the skin and deep fascia 5 min before the procedure. Facet RFT was applied first, followed by TFI and KE, respectively. RFT was applied for 60 s at 80°. Moreover, 10 mg bupivacaine (2 ml), 20 mg methylprednisolone (0.5 mL), and 0.9% isotonic sodium chloride solution (2.5 mL) combination for TFI and 40 mg bupivacaine (8 mL), 80 mg methylprednisolone (2 mL), and 0.9% isotonic sodium chloride solution (10 mL) combination for KE were applied. Iohexol 300 mg/50 mL was used as a radiopaque substance to determine the appropriate injection site in Els. Moreover, 3 mL of radiopaque fluid was used in TFI and 5 mL in CI. TFI and KE procedures were performed with Brauna® 22 gage (G), 90/120 mm Spinocan. Radiofrequency procedure was performed with TOP® 21 G, 100 mm, 10 mm

active cap, Neuropole Needle. Preganglionic epidural injection was performed via the subarticular route in the transforaminal procedure, and epidural injection was performed through the sacral hiatus in the caudal intervention. All procedures were performed under fluoroscopy guidance. Since it is known that successful results were obtained after only one treatment attempt in 94% of patients who underwent TFI, a single injection was attempted in all patients (20). After the procedure, the patients were transferred to the hospital bed and discharged after approximately 1 h of service follow-up, with oral nonsteroidal anti-inflammatory drugs prescribed.

**Statistical Analyses**

SPSS 25.0 (IBM Corporation, Armonk, NY) was used in the statistical analyses of the findings obtained in the study. Descriptive statistical methods (frequency, percentage, mean, standard deviation) were used to evaluate the study data in addition to Mann–Whitney U test and chi-squared test for the evaluation of normal distribution. The Mann–Whitney U test was used to compare the quantitative data between the two groups, the Wilcoxon sign test was used to compare whether there was a difference between the results of two measurements obtained from the same data source, and the Friedman test was used for measurements that were not in accordance with normal distribution. A p-value <0.05 was considered significant. Bonferroni correction was used to reduce the chances of obtaining false-positive results when multiple pair wise tests are performed on a single set of data.

**Table I:** Female and Male Ratios and Age Distribution in Relapse and Pathologies Other than Relapse

	Relapsed LDH (n=21)		Nonrelapsed LDH (n=27)	
	FB and TFI (n=8)	FB, TFI and CI (n=13)	FB and TFI (n=18)	FB, TFI and CI (n=9)
<b>Female</b>	3	5	11	5
<b>Male</b>	5	8	7	4
<b>Average age</b>	47 (27–64)	44 (24–73)	48 (30–69)	50 (37–80)

**FB:** Facet blockage, **TFI:** Transforaminal injection, **CI:** Caudal injection, **LDH:** Lumbar disc herniation.

**RESULTS**

The mean age of patients was 47.31 years (range, 24–80). There were 24 men (24–80 years; mean, 48.16 years) and 24 women (27–73 years; mean, 46.45 years) (age and female/male ratio, p=0.447 and p=0.140, respectively). Patients underwent LDS on an average of 66.41 months (1–300 months) before percutaneous interventions. There was no statistically significant difference in age (p=0.447) and sex (p=0.146) between both groups (Table I). In the radiological imaging of patients, 21 RDHs and 27 ODVP were detected (Table II).

Facet blockage (FB) and TFI were performed in 8 of RLDHs, and FB, TFI, and CI in 13. The mean age of those who underwent FB-TFI was 47 years (27–64), the F/M ratio was 3/5, the mean age of those who underwent FB-TFI-CI was 44 years (24–73), and the F/M ratio was 5/8. The mean age of the patients was 46 years (24–73), and the F/M ratio was 8/13. FB

**Table II:** Patient Characteristics and Comparison of Demographic Data

	Recurrence		p-value
	+(n=21)	-(n=27)	
<b>Age</b> <small>mean ± SD (min–max)</small>	45.52 ± 13.98 (24–73)	48.79 ± 12.48 (30–80)	<sup>1</sup> 0.447
<b>F/M</b> <small>n</small>	<b>Female</b>	8	16
	<b>Male</b>	13	11

<sup>1</sup>Mann–Whitney U test, <sup>2</sup>Chi-squared test, **n:** Number of patients, **SD:** Standard deviation.

and TFI were performed in 18 of ODVP, and FB, TFI and CI were performed in 9. The mean age of those who underwent FB-TFI was 48 years (30–69), the F/M ratio was 11/7, the mean age of those who underwent FB-TFI-CI was 50 years (37–80), and the F/M ratio was 5/4. The mean age of patients was 47.8 years (30–80), and the F/M ratio was 16/11. There was no statistically significant difference in age ( $p=0.447$ ) and sex ( $p=0.146$ ) between both groups (Table I).

Neurological examination results of the patients were followed closely by administering interventional pain treatments in the operating room environment under local anesthesia, and no permanent neurological deficits or major complications developed in the patients. However, severe pain was observed during the procedures in some patients. Nevertheless, none of the patients required discontinuation of interventional pain therapy.

The VAS and ODI scores of patients with RLDH who underwent FB-TFI were 8.62 and 42.62 preoperatively, 5.25

and 27.25 at 1 h postoperatively, and 4.33 and 25.42 at 6 months postoperatively, respectively. The VAS and ODI scores of patients with RLDH who underwent FB-TFI-CI were 8.61 and 42.84 preoperatively, 5 and 27.76 at 1 h postoperatively, and 4.88 and 28.22 at 6 months postoperatively, respectively (Table III).

The VAS and ODI scores of patients with ODVP who underwent FB-TFI were 8.55 and 44.5 preoperatively, 3.33 and 21.05 at 1 h postoperatively, and 4.11 and 24.11 at 6 months postoperatively, respectively. The VAS and ODI scores of patients with ODVP who underwent FB-TFI-CI were 8.55 and 41.55 preoperatively, 3.44 and 20 at 1 h postoperatively, 4.11 and 23 at 6 months postoperatively, respectively (Table III). In the follow-ups after interventional pain treatment, the ODI and VAS scores remained at a lower level compared to those preoperatively, and statistically significant improvement continued in the 6th month controls of patients with RLDH and ODVP who did not require surgery ( $p \leq 0.05$ ) (Tables IV, V).

**Table III:** VAS and ODI Scores Before, After, and 6 Months After Interventional Pain Treatment

	Relapsed LDH (n=21)		Nonrelapsed LDH (n=27)		
	FB and TFI ( $n=8$ )	FB, TFI and CI ( $n=13$ )	FB and TFI ( $n=18$ )	FB, TFI and CI (n=9)	
<b>Preoperative</b>	VAS	8.62 (8–9)	8.61 (8–9)	8.55 (7–9)	8.55 (7–9)
	ODI	42.62 (34–50)	42.84 (34–52)	44.5 (34–58)	41.55 (34–52)
<b>Postoperative</b>	VAS	5.25 (0–9)	5.0 (0–9)	3.33 (0–9)	3.44 (1–9)
	ODI	27.25 (10–46)	27.76 (11–42)	21.05 (6–58)	20 (10–40)
<b>Six months postoperative</b>	VAS	4.33 (1–9) <sup>δ</sup>	4.88 (1–9) <sup>δ</sup>	4.11 (0–9) <sup>δ</sup>	4.11 (1–9)
	ODI	25.42 (12–46) <sup>δ</sup>	28.22 (13–42) <sup>δ</sup>	24.11 (6–58) <sup>δ</sup>	23.00 (12–46)

<sup>δ</sup>Groups with patients who underwent operation in the follow-ups after interventional pain treatment. **LDH:** Lumbar disc herniation, **FB:** Facet blockage, **TFI:** Transforaminal injection, **CI:** Caudal injection.

**Table IV:** Baseline and Postprocedure Comparison of Oswestry Disability Index Scores Between the Groups

	Recurrence		p <sup>1</sup> -value
	+ (n=21)	- (n=27)	
Baseline ODI <sub>mean±SD (min-max)</sub>	42.76 ± 4.74 (34–52)	43.52 ± 6.11 (34–58)	0.867
<b>After the procedure</b>			
1 <sup>st</sup> hour ODI <sub>mean±SD (min-max)</sub>	27.57 ± 12.86 (10–46)	20.70 ± 15.14 (6–58)	0.055
6 <sup>th</sup> month ODI <sub>mean±SD (min-max)</sub>	34.52 ± 12.40 (12–50)	33.81 ± 17.24 (10–69)	0.892
P-value <sup>2</sup>	<0.001*	<0.001*	
<b>P-values between pairwise<sup>3</sup></b>			
Baseline vs. 1 <sup>st</sup> hour	0.002**	<0.001**	
Baseline vs. 6 <sup>th</sup> month	0.012**	0.009**	
1 <sup>st</sup> hour vs. 6 <sup>th</sup> month	0.006**	<0.001**	

Data presented as mean ± standard deviation (min-max). <sup>1</sup> Mann-Whitney U test <sup>2</sup>Friedman test <sup>3</sup>Wilcoxon signed-rank test  
**ODI:** Oswestry Disability Index. \*Statistically significant ( $p < 0.05$ ); \*\*Statistically significant ( $p < 0.05/3 = 0.017$ ) [according to Bonferroni correction]

**Table V:** Baseline and Postprocedure Comparison of Visual Analogue Scale Scores Between the Groups

	Recurrence		p-value <sup>1</sup>
	+(n=21)	-(n=27)	
Baseline VAS <small>mean ± SD (min-max)</small>	8.75 ± 0.45 (8-9)	8.58 ± 0.64 (7-9)	0.902
<b>After the procedure</b>			
1 <sup>st</sup> hour VAS <small>mean ± SD (min-max)</small>	4.56 ± 3.52 (0-9)	3.46 ± 2.97 (0-9)	0.136
6 <sup>th</sup> month VAS <small>mean ± SD (min-max)</small>	6.5 ± 8.03 (1-10)	4.12 ± 3.33 (0-9)	0.462
P-value <sup>2</sup>	<0.001*	<0.001*	
<b>P-values between pairwise</b>			
Baseline vs. 1 <sup>st</sup> hour	0.002**	<0.001**	
Baseline vs. 6 <sup>th</sup> month	0.049	<0.001**	
1st hour vs. 6 <sup>th</sup> month	0.058	0.035	

Data presented as mean ± standard deviation (min-max). <sup>1</sup> Mann-Whitney U test <sup>2</sup>Friedman test <sup>3</sup> Wilcoxon signed-rank test  
**VAS:** Visual analogue scale. \* Statistically significant ( $p < 0.05$ ); \*\* Statistically significant ( $p < 0.05/3 = 0.017$ ) [according to Bonferroni correction].

**Table VI:** Statistical Table of VAS and ODI Scores in Relapse and Nonrelapse Pathologies

	FB-TFI-CI (n)	FB-TFI (n)	p-value
<b>ODI</b>			
Relapse	9	16	0.284
Nonrelapse	7	10	1.000
<b>VAS</b>			
Relapse	8	9	0.248
Nonrelapse	13	15	0.386

**FB:** Facet blockage, **TFI:** Transforaminal injection, **CI:** Caudal injection, **ODI:** Oswestry Disability Index, **VAS:** Visual analog scale.

When the FB and TFI groups of these patients were compared with the groups treated with FB, TFI and CI, no statistically significant difference was found in the ODI and VAS scores at 6 months ( $p \geq 0.05$ ) (Table VI).

Among the patients with RLDH, 2 of 8 patients who underwent FB and TFI and 4 of 13 patients who underwent FB, TFI, and CI were operated within 6 months after the procedure. While 1 of 18 patients who underwent FB and TFI due to pathologies other than RLDH required surgical treatment within 6 months, no patient in the other group underwent surgical treatment. The numbers of patients whose complaints had decreased 50% and more than six months and 1 year in the RLDH group that underwent FB and TFI were 4/8 (50%) and 3/8 (37.5%), respectively, and in those who underwent FB, TFI and CI were 7/9 (77.7%) and 4/9 (44.4%), respectively. The success rates in patients treated the FB and TFI groups that applied to those with pathologies other than relapse were 11/18 (61.1%) and 8/18 (44.4%), respectively, and in those treated with FB, TFI, and CE were 7/9 (77.7%) and 4/9 (44.4%), respectively (Table VII, VIII).

The comparison between the reduction in pain level in the first hour after pain management procedures and continuation of well-being was investigated in all patients who had an operative history. In 85% (17/20) of patients whose pain level decreased by 71% or more according to the VAS scores measured at 1 hour after pain treatment, the benefit was sustained for 1 year or longer. These findings were statistically significant ( $p \leq 0.001$ ).

## ■ DISCUSSION

RLDHs and previously mentioned other pathologies response to the interventional pain treatment are different, which are the cause of recurrent LBP and leg pain after LDH surgery (17). There is no consensus on how to treat patients who underwent LDH surgery when similar complaints reappear. In clinical practice, most of the time, the treatments are administered based on clinical habits and experience relating to surgical or conservative treatments. At this point, these patients often do not have neurological deficit and that the purpose of treatment is to relieve pain. In the literature, the rate of complaints of low back and leg pain in the short-term (2 years ago) is 3%–34%, and the rate of similar complaints in the long-term (2 years later) is 5%–36% (26,33). At this point of view, starting from the most conservative treatment in the treatment of patients' pain to not perform unnecessary surgical interventions to the patients.

Surgical treatments of RLDs or other pathological conditions prementioned in the introduction have important complications, such as root injury, dural injury, increased fibrosis, and infections, and additional complications related to the general anesthesia (28). When complications related to TFIs are examined, spinal cord infarction has been reported in case reports (11,16,34). In the study reported of 14,956 TFIs' side effects, no complications that could cause serious damage, such as hematoma, infection, and permanent neurological injury, were reported, and minor and temporary complications,

**Table VII:** Patients Amount Whose Complaints had been Decreased %50 and More on Six Months and more than 1 Year

	Recurrent LDH		ODVP	
	FB-TFI	FB-TFI-CI	FB-TFI	FB-TFI-CI
Six months	4/8 (50%)	7/9 (77.7%)	11/18 (61.1%)	7/9 (77.7%)
>1 year	3/8 (37.5%)	4/9 (44.4%)	8/18 (44.4%)	4/9 (44.4%)

**FB:** Facet blockage, **TFI:** Transforaminal injection, **CI:** Caudal injection, **LDH:** lumbar disc herniation, **ODVP:** Other discovertebral pathologies.

**Table VIII:** Comparison of the Pain Rates in the Postoperative 1<sup>st</sup> Hour After the Procedure and the Duration

The rate of pain reduction at postoperative 1 <sup>st</sup> hour	Well-being period	
	1 year (19)	6 month (8)
71%–100% (n=20)	17 (89%)	1 (12.5%)
70%–61%	1 (5.5%)	2 (25%)
60%–51%	1 (5.5%)	4 (50%)
50%–41%	-	1 (12.5%)
40%–31%	-	-

such as vagovagal reaction, flushing, headache, insomnia, allergic reaction, changes in blood glucose level, and temporary increased weakness, were presented (6). In our study, no major life-threatening or neurological complications that cause permanent neurological deficits were observed in the patients.

Studies showed that TFIs applied to patients with primary LDH prevent the need for surgery (21,31). Although there is no such study on RLDHs, EIs may be effective in avoiding a new surgical treatment as much as possible for low back and leg pain after LDS and, in a sense, minimizing the risk of failure of back surgery. Since the patients included in the study had lower back pain in addition to leg pain, we also applied RFT in addition to EIs. The effectiveness of RFT has been demonstrated especially in LBP originating from the facet joint (4,6,18). However, corticosteroid injection into the facet joint was not performed with RFT, since it was shown that steroid injection together with RFT did not prevent neuritis formation after neurotomy and did not contribute to long-term results (1,30). The fact that facet RFT was performed on all patients also contributed to the standardization of the study. Moreover, in the study of Wei et al., the combined application of TFI and RFT had better long-term effectiveness and patient satisfaction, and that the recurrence rate was lower than the application of TFI alone (37).

In the study conducted by Klessinger, the success rate of TFI in ODVPs was 43%, and the relapse rate was 26.8%. It was stated that TFI may be effective in ODVPs, but the success rates in relapses should be further investigated (17). In our study, the success rate of TFI and FB performed in RLDHs were 50% in the 6<sup>th</sup> month, and 37.5% in the first year, while the success rate in ODVPs were 61.1% in the 6<sup>th</sup> month, and

44.4% in the first year. Although the steroid dose administered per level in the same study was 10 mg of triamcinolone, the administration of 20 mg of methylprednisolone per level may explain the higher success rates in our study. Apart from that, when the success rates in RLDHs and ODVPs were compared in our study, similarly better results were obtained (17). The study of Karamouzian et al., revealed that TFI and CI were successful in pain relief in recurrent disc hernias by approximately 40%, and there was no statistical difference between TFI and CI (14). They also emphasized that they could not achieve a high success rate in relapses (14). On the other hand, Evran et al. suggested that when patients who underwent TFI with TFI and CI were compared, those who underwent TFI and CI in RLDHs were statistically significantly better at the 3<sup>rd</sup> week, 3<sup>rd</sup> month, and 6<sup>th</sup> mo controls. It has been emphasized that the combined application of TFI and KE is more effective in RLDHs (7). In our study, no significant difference was found between the ODI and VAS scores of the patients in both groups who underwent FB+TFI and those who underwent FB+TFI+CI. We performed CI in patients with multiple disc herniation and in both groups. However, we did not see any particular difference to the CI in addition to the TFI. Thus, we suggested that EIs are more effective in ODVPs, and the rate of patients who need to surgical treatment is low.

When there was reduction in pain levels of  $\geq 71\%$  within the first h after the procedure in pain treatment applied to relapse and ODVPs, it has been determined that 85% of the cases remained well for more than 1 year. In the current literature, there isn't any study showed the relationship between reduction in pain level and maintenance of well-being. Additionally, in studies on EIs that were applied to RLDHs, there is no standard practice regarding the type and dose of steroids and local anesthetics (23,29). Moreover, although there are placebo studies on lumbar disc hernias that applied TFI, the lack of comparison with placebo in TFI studies on RLDHs raises doubts about steroid injection (35). Although the results are worse in RLDHs compared to other pathologies, since the long-term (>2 years) follow-up results cannot be evaluated, the benefits arising from the long-term effects of the injections cannot be evaluated, so it may not be appropriate to decide on the effectiveness by looking at the short-term results. To evaluate this, it was suggested that it would be beneficial to conduct studies with a long follow-up period. Another important point is that the power balances differ among the lumbar disc levels of the spine, which is a biomechanical structure (38). Since the biomechanical properties of the injected levels differ within themselves, perhaps evaluating the TFIs according to the levels may contribute to the formation of more accurate results.

Considering the biochemical properties of steroids and their effects on cells, it is a predictable fact that it will be effective when applied to the epidural area (12,32). However, to evaluate its effects in primary, relapse, and other pathologies, controlled studies with standard drug dose and type specific to patient groups and long follow-up periods are needed.

Although there are many articles in the literature about palliative pain procedures applied in spinal problems, there is almost no study on this subject in patient groups who have undergone LDS and whose pain complaints continue or recur. Although this study was conducted retrospectively and in a relatively small group of patients, we suggest that it is important in terms of the results obtained.

## ■ CONCLUSION

Interventional pain treatments were generally beneficial in patients with low back and leg pain after LDH surgery. We did not detect any differences in the aspect of effectiveness of pain procedures between recurrent disc herniation and other pathologies. We also determined that adding CI to the treatment protocol did not change the outcomes of pain management. However, considering the number of groups formed in the study, it would be appropriate to conduct the study with a larger patient series to obtain more reliable results.

### AUTHORSHIP CONTRIBUTION

Study conception and design: FD, SK, EC

Data collection: FD

Analysis and interpretation of results: FD, OO, YBO, SK, EC

Draft manuscript preparation: FD, OO, YBO, SK, EC

Critical revision of the article: FD, OO, YBO, SK, EC

Other (study supervision, fundings, materials, etc...): FD, OO, OB, YBO, SK, EC

All authors (FD, OO, OB, YBO, SK, EC) reviewed the results and approved the final version of the manuscript.

## ■ REFERENCES

1. Abd-Elseyed A, Loebertman M, Huynh P, Urits I, Viswanath O, Sehgal N: The long-term efficacy of radiofrequency ablation with and without steroid injection. *Psychopharmacology Bulletin* 50(4 Suppl 1):11-16, 2020
2. Aydin Y, Ziyal IM, Duman H, Turkmen CS, Basak M, Sahin Y: Clinical and radiological results of lumbar microdiscectomy technique with preserving of ligamentum flavum comparing to the standard microdiscectomy technique. *Surg Neurol* 57(1):5-13, 2002
3. Buchmann N, Preuß A, Gempt J, Ryang YM, Vazan M, Stoffel M, Meyer B, Ringel F: Outcome after surgical treatment for late recurrent lumbar disc herniations in standard open microsurgery. *World Neurosurg* 89:382-386, 2016
4. Civelek E, Cansever T, Kabatas S, Kircelli A, Yilmaz C, Musluman M, Ofluoglu D, Caner H: Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg* 22(2):200-206, 2012
5. Drazin D, Ugiliweneza B, Al-Khouja L, Yang D, Johnson P, Kim T, Boakye M: Treatment of recurrent disc herniation: A systematic review. *Cureus* 8(5):e622, 2016
6. El-Yahouchi CA, Plastaras CT, Maus TP, Carr CM, McCormick ZL, Geske JR, Smuck M, Pingree MJ, Kennedy DJ: Adverse event rates associated with transforaminal and interlaminar epidural steroid injections: A multi-institutional study. *Pain Med* 17(2):239-247, 2016
7. Evran S, Kayhan A, Baran O, Saygi T, Katar S, Akkaya E, Ozbek MA, Cevik S: The synergistic effect of combined transforaminal and caudal epidural steroid injection in recurrent lumbar disc herniations. *Cureus* 13(1): e12538, 2021
8. Fritsch EW, Heisel J, Rupp S: The failed back surgery syndrome: Reasons, intraoperative findings, and long-term results: A report of 182 operative treatments. *Spine* 21(5):626-633, 1996
9. Gaston P, Marshall RW: Survival analysis is a better estimate of recurrent disc herniation. *J Bone Joint Surg Br* 85(4):535-537, 2003
10. Ghahreman A, Bogduk N: Predictors of a favorable response to transforaminal injection of steroids in patients with lumbar radicular pain due to disc herniation. *Pain Med* 12(6):871-879, 2011
11. Gharibo CG, Fakhry M, Diwan S, Kaye AD: Conus medullaris infarction after a right L4 transforaminal epidural steroid injection using dexamethasone. *Pain Physician* 19(8):E1211-1214, 2016
12. Hassan KZ, Sherman AI: Epidural Steroids. [Updated 2022 Jul 19]. Treasure Island (FL): StatPearls Publishing, 2022
13. Hsu E, Atanelov L, Plunkett AR, Chai N, Chen Y, Cohen SP: Epidural lysis of adhesions for failed back surgery and spinal stenosis: Factors associated with treatment outcome. *Anesth Analg* 118(1):215-224, 2014
14. Karamouzian S, Ebrahimi-Nejad A, Shahsavarani S, Keikhosravi E, Shahba M, Ebrahimi F: Comparison of two methods of epidural steroid injection in the treatment of recurrent lumbar disc herniation. *Asian Spine J* 8(5):646-652, 2014
15. Karppinen J, Ohinmaa A, Malmivaara A, Kurunlahti M, Kyllönen E, Pienimäki T, Nieminen P, Tervonen O, Vanharanta H: Cost effectiveness of periradicular infiltration for sciatica: Subgroup analysis of a randomized controlled trial. *Spine* 26(23):2587-2595, 2001
16. Kennedy DJ, Dreyfuss P, Aprill CN, Bogduk N: Paraplegia following image-guided transforaminal lumbar spine epidural steroid injection: Two case reports. *Pain Med* 10(8):1389-1394, 2009
17. Klessinger S: Radicular pain in post lumbar surgery syndrome: The significance of transforaminal injection of steroids. *Pain Med* 14(2):243-246, 2013
18. Lee CH, Chung CK, Kim CH: The efficacy of conventional radiofrequency denervation in patients with chronic low back pain originating from the facet joints: A meta-analysis of randomized controlled trials. *Spine J* 17(11):1770-1780, 2017

19. Lee JH, Shin KH, Bahk SJ, Lee GJ, Kim DH, Lee CH, Kim DH, Yang HS, Lee SH: Comparison of clinical efficacy of transforaminal and caudal epidural steroid injection in lumbar and lumbosacral disc herniation: A systematic review and meta-analysis. *Spine J* 18(12):2343-2353, 2018
20. MacVicar J, King W, Landers MH, Bogduk N: The effectiveness of lumbar transforaminal injection of steroids: A comprehensive review with systematic analysis of the published data. *Pain Med* 14(1):14-28, 2013
21. Manson NA, McKeon MD, Abraham EP: Transforaminal epidural steroid injections prevent the need for surgery in patients with sciatica secondary to lumbar disc herniation: A retrospective case series. *Can J Surg* 56(2):89-96, 2013
22. Mascarenhas AA, Thomas I, Sharma G, Cherian JJ: Clinical and radiological instability following standard fenestration discectomy. *Indian J Orthop* 43(4):347-351, 2009
23. Mattie R, Miller DC, Smith C: Annual maximum dose of epidural steroid injection. *Pain Med* 20(10):2069-2070, 2019
24. Ozgen S, Naderi S, Ozek MM, Pamir MN: Findings and outcome of revision lumbar disc surgery. *J Spinal Disord* 12(4):287-292, 1999
25. Padua R, Padua S, Romanini E, Padua L, De Santis E: Ten- to 15-year outcome of surgery for lumbar disc herniation: Radiographic instability and clinical findings. *Eur Spine J* 8(1):70-74, 1999
26. Parker SL, Mendenhall SK, Godil SS, Sivasubramanian P, Cahill K, Ziewacz J, McGirt MJ: Incidence of low back pain after lumbar discectomy for herniated disc and its effect on patient-reported outcomes. *Clin Orthop Relat Res* 473(6):1988-1999, 2015
27. Shepard N, Cho W: Recurrent lumbar disc herniation: A review. *Global Spine J* 9(2):202-209, 2019
28. Shriver MF, Xie JJ, Tye EY, Rosenbaum BP, Kshetry VR, Benzel EC, Mroz TE: Lumbar microdiscectomy complication rates: A systematic review and meta-analysis. *Neurosurg Focus* 39(4):E6, 2015
29. Sim SE, Hong HJ, Roh K, Seo J, Moon HS: Relationship between epidural steroid dose and suppression of hypothalamus-pituitary-adrenal axis. *Pain Physician* 23(4S):S283-294, 2020
30. Singh JR, Miccio VF, Modi DJ, Sein MT: The impact of local steroid administration on the incidence of neuritis following lumbar facet radiofrequency neurotomy. *Pain Physician* 22(1):69-74, 2019
31. Smith CC, McCormick ZL, Mattie R, MacVicar J, Duszynski B, Stojanovic MP: The effectiveness of lumbar transforaminal injection of steroid for the treatment of radicular pain: A comprehensive review of the published data. *Pain Med* 21(3):472-487, 2020
32. Stafford MA, Peng P, Hill DA: Sciatica: A review of history, epidemiology, pathogenesis, and the role of epidural steroid injection in management. *Br J Anaesth* 99(4):461-473, 2007
33. Suri P, Pearson AM, Zhao W, Lurie JD, Scherer EA, Morgan TS, Weinstein JN: Pain recurrence after discectomy for symptomatic lumbar disc herniation. *Spine* 42(10):755-763, 2017
34. Tackla RD, Keller JT, Ernst RJ, Farley CW, Bohinski RJ: Conus medullaris syndrome after epidural steroid injection: Case report. *Int J Spine Surg* 6:29-33, 2012
35. Verheijen EJ, Bonke CA, Amorij EM, Vleggeert-Lankamp CL: Epidural steroid compared to placebo injection in sciatica: A systematic review and meta-analysis. *Eur Spine J* 30(11):3255-3264, 2021
36. Vik A, Zwart JA, Hulleberg G, Nygaard ØP: Eight year outcome after surgery for lumbar disc herniation: A comparison of reoperated and not reoperated patients. *Acta Neurochir* 143(6):607-611, 2001
37. Wei WB, Dang SJ, Wei L, Liu T, Wang J: Transforaminal epidural steroid injection combined with radio frequency for the treatment of lumbar disc herniation: A 2-year follow-up. *BMC Musculoskelet Disord* 22(1):347, 2021
38. Wilke HJ, Volkheimer D: Basic biomechanics of the lumbar spine. In: Galbusera F, Wilke HJ (eds), *Biomechanics of the Spine: Basic Concepts, Spinal Disorders and Treatments*. London: Elsevier, 2018:51-67