



DOI: 10.5137/1019-5149.JTN.20007-17.1

Received: 25.01.2017 / Accepted: 27.02.2017

Published Online: 17.03.2017

Original Investigation

Open-door Laminoplasty with Preservation of Muscle Attachments of C2 and C7 for Cervical Spondylotic Myelopathy: Retrospective Study

Halil Ibrahim SECER¹, Ferhat HARMAN², Murat Hamit AYTAZ³, Serdar KAHRAMAN⁴

¹University of Kyrenia, Faculty of Medicine, Department of Neurosurgery, Kyrenia, Turkish Republic of North Cyprus
Visiting Professor at Near East University, Faculty of Medicine, Nicosia, Turkish Republic of North Cyprus

²Marmara University, Faculty of Medicine, Department of Neurosurgery, Istanbul, Turkey

³Acibadem University, Vocational School of Health Services, Department of First Aid and Reanimation, Istanbul, Turkey

⁴Anadolu Medical Center, Department of Neurosurgery, Istanbul, Turkey

ABSTRACT

AIM: Some restriction and complications, such as progression of kyphosis, incidence of axial neck pain and decrease of postoperative cervical range of motion are concern. We designed this retrospective clinical study to evaluate the effect of laminoplasty by preserving the muscle attachments of C2 and C7 spinous processes on range of motion (ROM), axial neck pain and cervical lordosis.

MATERIAL and METHODS: Twenty-seven cases with cervical spondylotic myelopathy underwent open-door laminoplasty with the protection of muscle attachments to the C2 and C7 spinous process and laminae between 2007 and 2013. At the end of the follow-up, cases were evaluated with preoperative and postoperative modified Japanese Orthopedic Association (mJOA) scores, recovery rate, ROM, lordosis angle and visual analogue scale (VAS). Also, patients were divided into two groups and evaluated according to the magnetic resonance imaging (MRI) findings, with or without T2 signal change.

RESULTS: The mean age of the patients was 66 years. The mean follow-up duration was 25 months. The postoperative mJOA scores were significantly higher than the preoperative mJOA scores ($p < 0.001$). The recovery rate was 57.4%. Although the postoperative VAS score was higher than the preoperative VAS score and the mean postoperative ROM was lower than the preoperative ROM there was no significant difference between preoperative and postoperative VAS score and ROM ($p > 0.05$). The postoperative lordosis angle was significantly lower than the preoperative lordosis angle ($p < 0.05$). There were no significant differences regarding the postoperative lordosis angle, ROM and mJOA scores with or without T2 signal change on MRI.

CONCLUSION: Protection of the anatomic structures around the cervical spine such as the muscles and ligaments provides us better results regarding ROM and cervical axial pain.

KEYWORDS: Laminoplasty, Cervical spondylotic myelopathy, Range of motion, Lordosis

INTRODUCTION

Laminoplasty is a good alternative to laminectomy for the treatment of cervical spinal stenosis with/without myelopathy (3,18,20,21,36-38). The expansive laminoplasty technique was first described by Oyama et al. (26). After that, the unilateral open-door laminoplasty

technique was modified and developed by Hirabayashi and the double-door laminoplasty technique by Kurokawa (5,20). Although cervical laminoplasty is a motion-preserving procedure, several studies have reported some complications and problems such as axial neck pain, kyphotic deformity, and reduction of cervical range of motion (ROM) on long-term follow-up (6,8,20).

qr code

Corresponding author: Halil Ibrahim SECER

E-mail: hisecer@yahoo.com

In recent procedures, preservation of muscle attachments is a new and important development for the laminoplasty technique to reduce axial neck pain and restriction of cervical ROM. They all reach the consensus that protection of muscle attachments positively contributed to the reduction of pain and ROM (17,19,30-35,38).

Despite wide clinical experience about laminoplasty, the efficacy of and best procedure for laminoplasty remains unclear. We designed this retrospective study to evaluate the patients who underwent open-door laminoplasty with preservation of muscle attachments of C2 and C7 spinous processes for cervical spondylotic myelopathy and compared the preoperative and postoperative clinical and radiological features.

■ MATERIAL and METHODS

Patient Demographics

There were 27 consecutive patients with cervical spondylotic myelopathy. All the patients underwent the open-door laminoplasty procedure by two experienced surgeons between December 2007 and October 2013. There were 20 males (74%) and 7 females (26%). The mean age at the time of surgery was 66 years (range 45-83 years). The mean duration of the symptoms before the surgery was 14 months (range 3-42 months). The mean follow-up duration was 25 months (range 6-45 months).

At the end of the follow-up period, each patient's status was evaluated with preoperative and postoperative modified Japanese Orthopedic Association (mJOA) scores, recovery rate, ROM, lordosis angle and visual analogue scale (VAS). Also, the patients were divided into two groups and evaluated according to the magnetic resonance imaging (MRI) findings, with or without T2 signal change. There were 12 patients who had a T2 signal change in the spinal cord on MRI scan (T2-MRI (+) group) and 15 patients who did not have a signal change in the spinal cord on MRI scan (T2-MRI (-) group).

Surgical Procedures

The head was fixed with a Mayfield frame and the patient was rotated to the prone position. A posterior midline C2-C7 skin incision was made. The paravertebral muscles were dissected from the spinous processes and lamina to the facet joints, but facet capsules were preserved to avoid fusion. Muscles attached to the C2 and C7 spinous process and laminae were all protected. The supraspinous and interspinous ligaments were strictly preserved. The patient's more painful or neurologically affected side was chosen as the laminotomy side. The operating microscope was routinely used during all laminoplasty procedures. A triangle-shaped bone gutter at the medial margin of the facet joints was made at the contralateral side with a high-speed drill. The laminotomy was performed with a high-speed drill or a Number 1 Kerrison rongeur at one side. The lamina and ligamentum flavum were elevated from the open side and hinged on the other side, which decompressed and relieved the spinal cord. A titanium laminoplasty plate was used to maintain patency of the

open-door laminoplasty (NewBridge laminoplasty fixation systems, Orthofix, Lewisville, USA). Foraminotomy was performed in cases of myeloradiculopathy. C5 foraminotomy was not performed routinely. Postoperatively, the patients wore a soft cervical collar for 2 weeks.

Neurological Assessment

Postoperative results were evaluated by using the mJOA score (14). Using a formula, the degree of postoperative neurological recovery was calculated as follows (5).

Recovery rate (%) = (postoperative mJOA score - preoperative mJOA score) / (full mJOA score [18] - preoperative mJOA score) x 100.

Radiological Assessments

Preoperative and postoperative ROM, and the preoperative and postoperative lordosis angle were measured on the lateral cervical spine x-rays according to the relevant literature. The angle of cervical lordosis angle was measured between a line parallel to the posterior surface of the body of the C2 and a line parallel to the posterior surface of the body of C7 (13).

Statistical Analyses

Statistical analyses were performed using the IBM SPSS for Windows Version 21.0 statistical package. Continuous variables were presented as mean ± standard [minimum - maximum]. Categorical variables were summarized as frequencies and percentages. Normality of the continuous variables was evaluated by Shapiro-Wilk test. Differences between the two groups according to continuous variables were determined by Mann-Whitney U test. Differences between preoperative and postoperative values were evaluated by the Wilcoxon signed rank test. A p value less than 0.05 was considered as significant.

■ RESULTS

In all patients, the mean operative time was 105 minutes (range 90-150 minutes), and the mean estimated blood loss was 355 ml (range 200-600 ml). The mean age was 66.8 ± 8.2 years (range 45-83 years). Laminoplasty was started from the C3 level in 18 cases (66%). Laminoplasty was performed on 4 levels in 13 cases, 3 levels in 12 cases and 2 levels in only 2 cases. There were T2 signal changes in 12 cases (Table I).

The mean preoperative mJOA scores were 12.2 ± 2.4 and postoperative mJOA scores were 15.0 ± 2.5. The postoperative mJOA scores were significantly higher than the preoperative mJOA scores (p < 0.001). The recovery rate was 57.4% (Table II).

The mean preoperative VAS score was 5.6 ± 1.4 and the postoperative VAS score was 6.2 ± 1.9. Although the postoperative VAS score was higher than the preoperative VAS score, there was no statistically significant difference between them (p > 0.05) (Table II).

The postoperative ROM decreased in the 22 cases, did not change in 2 cases and increased in 3 cases when compared with the preoperative ROM. The mean preoperative ROM

was 35.3 ± 5.3 (range 22-43) degrees and postoperative ROM was 34.0 ± 6.4 (range 20-52) degrees. Although the mean postoperative ROM was lower than the preoperative ROM,

Table I: Patient Characteristics

	Mean±SD	
Age	66.8±8.2 years	
Duration of Symptoms	14.0±9.3 months	
Recovery rate	57.4±28.3	
Duration of Follow-up	25.0±10.7 months	
	Frequency	
MRI	-	15
	T2	12
Level	2 (C4-C5)	1
	2 (C5-C6)	1
	3 (C3-C5)	5
	3 (C4-C6)	7
	4 (C3-C6)	13
	2	2
Number of Level	3	12
	4	13
C5 Palsy	-	24
	+	3

there was no statistically difference between them ($p > 0.05$) (Table II).

The preoperative cervical lordosis angle ranged from -10 to 37 degrees and cases with cervical kyphosis higher than -10 degrees were not operated on with this techniques. Postoperative cervical lordosis angles ranged from -15 to 30 degrees and decreased in 16 cases, did not change in 8 cases and increased in 3 cases when compared with the preoperative lordosis angle. The mean preoperative lordosis angle was $14.1^\circ \pm 13.5^\circ$ (range $-10^\circ \pm 37^\circ$) and postoperative lordosis angle was $12.6^\circ \pm 13.5^\circ$ (range $-15^\circ \pm 30^\circ$). The postoperative lordosis angle was significantly lower than the preoperative lordosis angle ($p < 0.05$) (Table II).

While evaluating the recovery rate according to the T2 signal change in the spinal cord on MRI scan, the mean recovery rate was 73.6 ± 25.2 in the T2-MRI (+) group and 37.1 ± 16.8 in the T2-MRI (-) group. The recovery rate of the T2-MRI (+) group was significantly higher than the T2-MRI (-) group ($p < 0.05$). There were no significant differences regarding the postoperative lordosis angle, ROM and mJOA scores between these two groups (Table III).

Temporary C5 nerve root palsy had occurred on the hinged side of three patients (11.1%). All patients with C5 nerve root palsy were completely recovered within 3 months. In one patient, deep surgical site infection developed. *Staphylococcus aureus* was isolated from the culture material and the condition cured within six weeks using vancomycin (Kocak Farma Pharmacy and Chemical Industry, Istanbul, Turkey) treatment. One patient had a C6 screw pull-out at the second month of operation and was revised surgically. While elevating the lamina, the hinged side was broken in 2 patients and fixed with a miniplate/screw system.

Table II: The Results and Statistical Analysis of Lordosis and ROM Degrees, JOA and VAS Scores

	Preoperative	Postoperative	p
	Mean±SD	Mean±SD	
Lordosis degree	14.1 ± 13.5	12.6 ± 13.5	0.005
mJOA score	12.2 ± 2.4	15.0 ± 2.5	<0.001
ROM	35.3 ± 5.3	34.0 ± 6.4	0.216
VAS	5.6 ± 1.4	6.2 ± 1.9	0.790

Table III: Analyzing of the Effect of MRI T2 Signal Change in the Spinal Cord on Recovery Rate, mJOA Score, Lordosis and ROM Degrees

	MRI (-)	MRI (T2+)	p
	Mean±SD	Mean±SD	
Recovery rate	73.6 ± 25.2	37.1 ± 16.8	0.001
Δ Lordosis degree	3.5 ± 3.8	-0.9 ± 8.1	0.139
Δ mJOA score	-3.0 ± 1.4	-2.7 ± 1.1	0.755
Δ ROM	1.1 ± 4.2	2.6 ± 2.0	0.399

■ DISCUSSION

During the past decades, laminoplasty has become a good treatment choice for many surgeons in cases of cervical spondylotic myelopathy after the development of the open-door laminoplasty technique. There are two different mechanisms ensuring decompression of the spinal cord with the laminoplasty method. Firstly, direct local decompression of the posterior elements decreases the posterior pressure on the spinal cord. Secondly, the dorsal shift of the spinal cord moves it away from the anterior compressing factors such as osteophytes, ossified-hypertrophic ligaments and protruding discs, obtaining a total decompression effect (7).

Many modifications such as bone grafts, hydroxyapatite or other ceramic materials, allograft and titanium plates, titanium miniplates and screws have been developed to prevent this complication (2,7,20,21,39). We have preferred titanium miniplates (Blackstone Medical Inc, USA). We achieved good results with this technique. While elevating the lamina, the hinged side was broken in only 2 patients and fixed with a miniplate/screw system. Just one patient had a C6 screw pull-out at the second month of operation and was revised surgically.

In the earliest Hirabayashi report, open-door laminoplasty was performed in 40 patients with multilevel cervical spondylotic myelopathy and most of the patients had good results. The recovery rate was 66% with the JOA scale score in this series. Many other surgeons reported similar results with JOA scores and neurological recovery rates after various techniques of laminoplasty. They reported the improvement to be stable for more than 10 years in most patients (1,22,29). We found that the postoperative mJOA scores were significantly higher than the preoperative mJOA scores and the recovery rate was 57.4% in our study which is similar to the previous literature.

Although cervical laminoplasty is an effective decompressing and motion-preserving procedure, several studies have reported some complications and problems such as axial neck pain, kyphotic deformity, and reduction of cervical ROM on long-term follow-up (6,8,20). In this study, although the postoperative lordosis degree was statistically lower than the preoperative lordosis degree, there was no statistically significant difference between the postoperative ROM and the preoperative ROM. Kang et al. retrospectively evaluated 20 laminoplasty patients with their preoperative and postoperative ROM and they determined factors (JOA score, duration of symptoms, disease entity, age and sex) associated with cervical ROM. They found cervical ROM was reduced after laminoplasty and there was no correlative factor that was associated with a reduction in cervical ROM (16). Duetzmann et al. presented a wide literature review about 103 laminoplasty series and noted that in all series a mean 47.3% loss of ROM was reported (2). We found a decrease in neck flexibility after the laminoplasty procedure, but there was no statistically significant difference.

Kyphosis can develop after cervical laminoplasty in the straight or lordotic cervical spine. Iwasaki et al. reported as 8% progressive kyphotic deformity after cervical laminoplasty,

but no neurological deterioration (11). Some authors presented that detachment of cervical muscles, especially the semispinalis cervicis muscle attached to the C2 spina, were associated with postoperative kyphosis (12,20). No detachment and preservation of the deep extensor muscles to the C2 spinous process was correlated with postoperative changes in cervical kyphotic deterioration (2,12,32-34,38). Duetzmann et al. reviewed a total of 103 studies from MEDLINE. They showed that only 21 studies had statistical comparisons between preoperative and postoperative C2-C7 angles. Moreover, 28.8% of these studies reported had stable alignment, 45.0% had worse kyphosis, and 26.3% had increased lordotic curvature. They emphasized that understanding the muscle-ligament complex is important for better surgical results of cervical alignment. In our cases, the muscles attached to the C2 and C7 spinous process and the supraspinous and interspinous ligaments were strictly preserved preoperatively (2).

Hardacker et al. and Scheer et al. evaluated the cervical sagittal parameters and they found cervical lordosis angle was not occurred evident symptom until that angle decreased to -9.6 degrees (4,28). We did not operate on patients with a lordosis angle lower than -10 degrees. However, in three cases, the postoperative lordosis angle increased to over -10 degrees when compared with the preoperative lordosis angle (-8 to -12, -5 to -12, and -10 to -15 degree). In one case, the preoperative lordosis angle increased from -10 to +16 degrees. In this study, the postoperative lordosis angle was significantly lower than the preoperative lordosis angle. Protection of the C2 and C7 spinous-attached muscles during the surgery and not operating on patients with cervical lordosis lower than -10 degrees is important for maintenance of cervical lordosis after laminoplasty. Even if ROM and axial neck pain had negative effects, it did not cause a statistically significant change.

Many authors have reported axial neck pain in 27%–90% of patients after laminoplasty for cervical spondylotic myelopathy (8,10,23). Causes of axial pain are still uncertain. Many studies have shown that preservation of C2 and C7 spinous processes reduced the frequency of postoperative axial neck pain (9,10,25,27,33,34,38). In the present study, the mean preoperative VAS score for axial neck pain was 5.6 ± 1.4 and the postoperative VAS score was 6.2 ± 1.9 at the end of the follow-up period. Although the postoperative VAS score was higher than the preoperative VAS score, there was no statistically significant difference between them ($p > 0.05$). One reason for the relative infrequency of axial neck pain among our patients was the preservation of the semispinalis muscle insertion into C2, and the paravertebral muscles at the C7. Another factor influencing axial neck pain is the use of a cervical soft collar in the early postoperative period. This should be shortened. In the present study, the brace therapy period was limited to 2 weeks, and active sitting, standing, and ambulation were started on the postoperative first day.

The C5 nerve root is the most commonly affected nerve root after open-door laminoplasty and nerve injury usually results in motor weakness (6,15). The tethering of the nerve roots with posterior migration of the spinal cord may cause this

problem. Patients with ossified posterior longitudinal ligament (OPLL) also have a higher risk of C5 nerve palsy (36). Gentle elevation of the lamina may be effective for the prevention of this complication (15). In our series, C5 nerve palsy occurred in 3 patients (11.1%) in the early postoperative period and all of them recovered within a few months. All of them were operated on four levels (C3-C6) and two had MRI T2 changes preoperatively.

Spinal cord signal changes on MRI of cervical spondylotic myelopathy patients have been correlated with preoperative and postoperative neurological condition and JOA scores. Sun et al. showed that T2-WI hyperintense MRI changes were present in approximately 80.7% of all patients and MRI parameters were significant predictors of surgical outcome (31). Nouri et al. prospectively evaluated 102 patient's MRI parameters and mJOA scores. They showed that MRI changes have a significant role in predicting the surgical outcome (24). Similarly in our study, we also evaluated T2-WI changes on MRI. Recovery rate of the T2-MRI (-) group was significantly higher than the T2-MRI (+) group. Change of lordosis angle, mJOA scores and ROM was not evident postoperatively between two groups.

■ CONCLUSION

Laminoplasty has become an effective treatment method for many cases of cervical spondylotic myelopathy after the development of the open-door laminoplasty technique. The patients with preoperative MRI changes in the spinal cord have worse post-laminoplasty outcome and neurological condition. Protection of the anatomical structures around the cervical spine such as the muscles attached to the C2 and C7 spinous process, and the supraspinous and interspinous ligaments provides better results regarding ROM and cervical axial pain.

■ REFERENCES

- Chiba K, Ogawa Y, Ishii K, Takaishi H, Nakamura M, Maruiwa H, Matsumoto M, Toyama Y: Long-term results of expansive open-door laminoplasty for cervical myelopathy—Average 14-year follow-up study. *Spine* 31: 2998–3005, 2006
- Duetzmann S, Cole T, Ratliff JK: Cervical laminoplasty developments and trends, 2003-2013: A systematic review. *J Neurosurg Spine* 23(1): 24-34, 2015
- Hale JJ, Gruson KI, Spivak JM: Laminoplasty: A review of its role in compressive cervical myelopathy. *Spine J* 6 Suppl 6: 289-298, 2006
- Hardacker JW, Shuford RF, Capicotto PN, Pryor PW: Radiographic standing cervical segmental alignment in adult volunteers without neck symptoms. *Spine* 22(13): 1472-1480, 1997
- Hirabayashi K, Satomi K: Operative procedure and results of expansive open-door laminoplasty. *Spine* 13(7): 870-876, 1988
- Hirabayashi K, Toyama Y, Chiba K: Expansive laminoplasty for myelopathy in ossification of the longitudinal ligament. *Clin Orthop Relat Res* 359: 35-48, 1999
- Hirabayashi K, Watanabe K, Wakano K, Suzuki N, Satomi K, Ishii Y: Expansive open-door laminoplasty for cervical spinal stenotic myelopathy. *Spine* 8(7): 693-699, 1983
- Hosono N, Sakaura H, Mukai Y, Fujii R, Yoshikawa H: C3-6 laminoplasty takes over C3-7 laminoplasty with significantly lower incidence of axial neck pain. *Eur Spine J* 15:1375–1379, 2006
- Hosono N, Sakaura H, Mukai Y, Ishii T, Yoshikawa H: En-bloc laminoplasty without dissection of paraspinal muscles. *J Neurosurg Spine* 3: 29–33, 2005
- Hosono N, Yonenobu K, Ono K: Neck and shoulder pain after laminoplasty. A noticeable complication. *Spine* 21(17):1969-1973, 1996
- Iwasaki M, Kawaguchi Y, Kimura T, Yonenobu K: Long-term results of expansive laminoplasty for ossification of the posterior longitudinal ligament of the cervical spine: More than 10 years follow up. *J Neurosurg* 96 Suppl 2: 180-189, 2002
- Izuka H, Shimizu T, Tateno K, Toda N, Edakuni H, Shimada H, Takagishi K: Extensor musculature of the cervical spine after laminoplasty: Morphologic evaluation by coronal view of the magnetic resonance image. *Spine* 26: 2220-2226, 2001
- Jackson R: *The Cervical Syndrome*, 2nd ed. Springfield, IL: Charles C. Thomas, 1958:174
- Japanese Orthopedic Association. Scoring system for cervical myelopathy. *J Jpn Orthop Assoc* 68: 490–503, 1994
- Kaminsky SB, Clark CR, Traynelis VC: Operative treatment of cervical spondylotic myelopathy and radiculopathy. A comparison of laminectomy and laminoplasty at five year average follow-up. *Iowa Orthop J* 24: 95-105, 2004
- Kang SH, Rhim SC, Roh SW, Jeon SR, Baek HC: Postlaminoplasty cervical range of motion: Early results. *J Neurosurg Spine* 6(5): 386-390, 2007
- Kim P, Murata H, Kurokawa R, Takaishi Y, Asakuno K, Kawamoto T: Myoarchitectonic spinolaminoplasty: Efficacy in reconstituting the cervical musculature and preserving biomechanical function. *J Neurosurg Spine* 7: 293-304, 2007
- Kimura A, Seichi A, Inoue H, Hoshino Y: Long-term results of double-door laminoplasty using hydroxyapatite spacers in patients with compressive cervical myelopathy. *Eur Spine J* 20: 1560–1566, 2011
- Kotani Y, Abumi K, Ito M, Sudo H, Takahata M, Ohshima S, Hojo Y, Minami A: Minimum 2-year outcome of cervical laminoplasty with deep extensor muscle-preserving approach: Impact on cervical spine function and quality of life. *Eur Spine J* 18(5): 663-671, 2009
- Kurokawa R, Kim P: Cervical laminoplasty: The history and the future. *Neurol Med Chir (Tokyo)* 55(7): 529-539, 2015
- Lin S, Zhou F, Sun Y, Chen Z, Zhang F, Pan S: The severity of operative invasion to the posterior muscular-ligament complex influences cervical sagittal balance after open-door laminoplasty. *Eur Spine J* 24(1): 127-135, 2015
- Machino M, Yukawa Y, Hida T, Ito K, Nakashima H, Kanbara S, Morita D, Kato F: Modified double-door laminoplasty in managing multilevel cervical spondylotic myelopathy: Surgical outcome in 520 patients and technique description. *J Spinal Disord Tech* 26: 135–140, 2013

23. Nakano N, Nakano T, Nakano K: Comparison of the results of laminectomy and open-door laminoplasty for cervical spondylotic myeloradiculopathy and ossification of the posterior longitudinal ligament. *Spine* 13: 792-794, 1988
24. Nouri A, Tetreault L, Zamorano JJ, Dalzell K, Davis AM, Mikulis D, Yee A, Fehlings MG: Role of magnetic resonance imaging in predicting surgical outcome in patients with cervical spondylotic myelopathy. *Spine* 40(3): 171-178, 2015
25. Ono A, Tonosaki Y, Yokoyama T, Aburakawa S, Takeuchi K, Numasawa T, Wada K, Kachi T, Toh S: Surgical anatomy of the nuchal muscles in the posterior cervicothoracic junction: Significance of the preservation of the C7 spinous process in cervical laminoplasty. *Spine* 33: E349-E354, 2008
26. Oyama M, Hattori S, Moriwaki N: A new method of cervical laminoplasty. *Centr Jpn J Orthop Traumatic Surg* 16: 792-794, 1973
27. Sakaura H, Hosono N, Mukai Y, Iwasaki M, Yoshikawa H: Medium-term outcomes of C3-6 laminoplasty for cervical myelopathy: A prospective study with a minimum 5-year follow-up. *Eur Spine J* 20: 928-933, 2011
28. Scheer JK, Tang JA, Smith JS, Acosta FL Jr, Protopsaltis TS, Blondel B, Bess S, Shaffrey CI, Deviren V, Lafage V, Schwab F, Ames CP; International Spine Study Group: Cervical spine alignment, sagittal deformity, and clinical implications: A review. *J Neurosurg Spine* 19(2): 141-159, 2013
29. Seichi A, Takeshita K, Ohishi I, Kawaguchi H, Akune T, Anamizu Y, Kitagawa T, Nakamura K: Long-term results of double-door laminoplasty for cervical stenotic myelopathy. *Spine* 26: 479-487, 2001
30. Shiraishi T: A new technique for exposure of the cervical spine laminae. Technical note. *J Neurosurg* 96: 122-126, 2002
31. Sun Y, Zhang F, Wang S, Zhang L, Pan S, Yu M, Qiu S: Open door expansive laminoplasty and postoperative axial symptoms: A comparative study between two different procedures. *Evid Based Spine Care J* 1(3): 27-33, 2010
32. Takeuchi K, Yokoyama T, Ono A, Numasawa T, Wada K, Kumagai G, Ito J, Ueyama K, Toh S: Cervical range of motion and alignment after laminoplasty preserving or reattaching the semispinalis cervicis inserted into axis. *J Spinal Disord Tech* 20: 571-576, 2007
33. Takeuchi K, Yokoyama T, Aburakawa S, Saito A, Numasawa T, Iwasaki T, Itabashi T, Okada A, Ito J, Ueyama K, Toh S: Axial symptoms after cervical laminoplasty with C3 laminectomy compared with conventional C3-C7 laminoplasty: A modified laminoplasty preserving the semispinalis cervicis inserted into axis. *Spine* 30: 2544-2549, 2005
34. Takeuchi K, Yokoyama T, Ono A, Numasawa T, Wada K, Itabashi T, Toh S: Limitation of activities of daily living accompanying reduced neck mobility after laminoplasty preserving or reattaching the semispinalis cervicis into axis. *Eur Spine J* 17(3): 415-420, 2008
35. Takeuchi T, Shono Y: Importance of preserving the C7 spinous process and attached nuchal ligament in French-door laminoplasty to reduce postoperative axial symptoms. *Eur Spine J* 16(9): 1417-1422, 2007
36. Tetreault L, Ibrahim A, Côté P, Singh A, Fehlings MG: A systemic review of clinical and surgical predictors of complications following surgery for degenerative cervical myelopathy. *J Neurosurg Spine* 25:1-23, 2015
37. Tomita K, Kawahara N, Toribatake Y, Heller JG: Expansive midline T-saw laminoplasty (modified spinous process-splitting) for the management of cervical myelopathy. *Spine* 23: 32-37, 1998
38. Tümtürk A, Kuçuk A, Menku A, Koc RK: En bloc cervical laminoplasty with preserving posterior structure and arcocristectomy in cervical spondylotic myelopathy. *Turk Neurosurg*, 2016 (Epub ahead of print)
39. Yeh KT, Yu TC, Chen IH, Peng CH, Liu KL, Lee RP, Wu WT: Expansive open-door laminoplasty secured with titanium miniplates is a good surgical method for multiple-level cervical stenosis. *J Orthop Surg Res* 9:49, 2014