



Posterior Cervical Discectomy via Keyhole Foraminotomy: A Case Series Comparing Microdiscectomy with Endoscope-Assisted Discectomy

Ali DALGIC¹, Resul KARADENİZ², Derya KARAOĞLU GUNDOĞDU³, Egemen ISITAN⁴, Goksal GUNERHAN², Ahmet Deniz BELEN²

¹Ankara Medica International Hospital, Neurosurgery Clinic, Ankara, Turkey

²Ankara City Hospital, Department of Neurosurgery, Ankara, Turkey

³Selcuk University, School of Medicine, Department of Neurosurgery, Konya, Turkey

⁴Hitit University, Erol Olçok Research and Training Hospital, Department of Neurosurgery, Corum, Turkey

Corresponding author: Ali DALGIC ✉ alidalgic@yahoo.com



To watch the surgical videoclip, please visit <http://turkishneurosurgery.org.tr/uploads/jtn-34918-video-1.mp4>
<http://turkishneurosurgery.org.tr/uploads/jtn-34918-video-2.mp4>

ABSTRACT

AIM: Anterior cervical discectomy and fusion (ACDF) has become the standard surgical treatment of cervical disc hernias (CDH), however, its complications including dysphagia, recurrent laryngeal nerve injury, pseudoarthrosis, implants and graft problems, adjacent segment disease have been well-known. Posterior cervical discectomy (PCD) via posterior keyhole lamino-foraminotomy may be a solution to avoid from these complications. To report the clinical outcomes of microdiscectomy (MD) and endoscope assisted discectomy (EAD) techniques via the posterior approach in patients with cervical disc herniations (CDHs).

MATERIAL and METHODS: The data were obtained from retrospective review of the patient's charts and the latest follow-up examination.

RESULTS: A total of 83 cases with CDH who were treated by posterior cervical discectomy (PCD), between 2010 and 2019, were reviewed. MD was used in 42 patients (male: 20, female: 22) with a mean age of 51.1 years. In MD group, all patients had pain, and 26 of them had additional weakness. Visual analogue scale (VAS) neck score was 7.72, VAS arm score was 8.83; PROLO score was 7.41. EAD was used in 41 patients (male: 26, female: 15) with a mean age of 38.7 years. In EAD group, all patients had pain, and 20 of them had additional weakness. VAS neck and arm scores were 7.75, and 8.72, respectively; PROLO score was 7.44. Mean follow-up time was 24.7 months. The scores at the latest exam are as follows: in MD group, VAS score for neck was 2.32, for arm 1.11; PROLO score was 9.58; in EAD group, VAS score for neck was 2.18 and for arm 0.97; PROLO score was 9.66. Both surgical techniques were success with statically significance ($p \leq 0.05$) according to the scores. The techniques were equally effective while postoperative VAS ($p > 0.412$) and PROLO ($p > 0.980$) scores were similar in both groups.

CONCLUSION: Both approaches are effective for selected patients with soft cervical disc herniation in which settled lateral location. Both techniques allow working with two handle, therefore facilitating the gentle manipulation that can obtain for avoiding hazardous effect to spinal cord and nerve root.

KEYWORDS: Cervical disc herniation, Key-hole foraminotomy, Endoscopy, Microdiscectomy, Posterior approach

ABBREVIATIONS: **ACDF:** Anterior cervical discectomy and fusion, **CDH:** Cervical disc hernia, **CT:** Computed tomography, **HO:** Heterotopic ossification, **MRI:** Magnetic resonance imaging, **PCD:** Posterior cervical discectomy, **PCF:** Posterior cervical foraminotomy, **PC-EAD:** Posterior cervical endoscope-assisted discectomy, **PC-MD:** Posterior cervical microdiscectomy, **PLL:** Posterior longitudinal ligament

Ali DALGIC : 0000-0003-1000-2811
Resul KARADENİZ : 0000-0001-7739-5619
Derya KARAOĞLU GUNDOĞDU : 0000-0003-2345-8818

Egemen ISITAN : 0000-0003-1993-050X
Goksal GUNERHAN : 0000-0001-6255-8315
Ahmet Deniz BELEN : 0000-0001-8863-9121

INTRODUCTION

Anterior cervical discectomy and fusion (ACDF) has become the standard surgical treatment of cervical disc hernias (38). Its complications include dysphagia, recurrent laryngeal nerve injury, and perforation of the esophagus (4,27,37,48). Pseudoarthrosis is the most common problem encountered during the follow-up period that could be addressed using implants and graft problems may require graft revision (29,32,44). Nevertheless, bone fusion was achieved despite the fact that adjacent segment disease may occur as the next trouble (8). Spine surgeons are working on means to prevent ACDF-related complications and posterior cervical discectomy (PCD) via posterior keyhole laminoforaminotomy is one of these milestone procedures (8,10,26). However, PCD requires muscle dissection reason why its use is not widespread. Recently, the introduction of the surgical microscope and speculum- or tube-like retractors minimizing muscular dissection has facilitated the posterior approach (7,12). Besides, PCD has been shown to shorten hospital stay, lessen tissue injury, and lower the amount of blood loss due to ACDF (14,47). Moreover, some reports reveal that PCD is an effective technique for preserving motion at the operation site (since it does not require fusion) and allows for early recovery, thus lowering the risk of adjacent segmental disease (24,28). Therefore, we retrospectively reviewed the postoperative clinical outcomes of PCD performed with posterior cervical microdiscectomy (PC-MD) or posterior cervical endoscope-assisted discectomy (PC-EAD) techniques.

MATERIAL and METHODS

Patients

From August 2010 to November 2019, 351 patients who suffered from cervical disc hernia (CDH) were treated at our hospital. PCD was performed in 104 patients of whom 83 were followed up. We reviewed the records of the latest control examinations of 42 (75%) patients who had PC-MD and 41 (85%) patients who had PC-EAD. We excluded patients whose follow-up period was shorter than 12 months.

A lateral cervical plain radiograph, a computed tomography (CT) scan, and a magnetic resonance imaging (MRI) scan were used to diagnose CDH, assess cervical spinal alignment, and to demonstrate degenerative changes (loss of height, black disc, calcification, osteophyte formation) in all patients. All patients underwent PC-MD or PC-EAD technique via a posterior approach to treat single-level, soft cervical disc herniation. Patients were included if they had signs of radiculopathy compatible with signs on a preoperative cervical MRI (Figure 1A-C), a laterally-located single-level soft disc hernia on radiological imaging, and those with persistent signs after conservative treatment for minimum four weeks, or whose symptoms worsened over this period. All patients were operated upon by the same surgeon (AD). The PC-MD technique was preferred in patients with soft disc accompanied by signs of degeneration (osteophytes) at a foraminal level on a preoperative plain film and/or CT scan, reduced disc height, and facet joint hypertrophy. PC-EAD was performed in patients without any sign of degeneration.




Figure 1: MRI scans revealed a soft disc herniation that settled at the left foramina of C6-7 (A), CT images showed degenerative findings such as reduced disc height, osteophytes at the different levels, but herniation was not calcified and no stenosis at the foramen (B), removed disc and effective decompression was seen postoperative MRI scans (C).

We excluded patients with segmental instability, CDH with midline location, posterior longitudinal ligament (PLL) calcification, central cervical stenosis and/or foraminal stenosis without CDH, multilevel CDH, or suspected infection or tumor in the cervical vertebrae. Patients who were followed up for less than 12 months were also excluded.

Surgical Technique

PC-MD; The patient was placed on the operating table in the prone position with their neck slightly flexed under general anesthesia. The disc level was determined using fluoroscopy. A skin incision of approximately 1.5 cm long was made in a linear fashion approximately 2 cm away from the midline to center of the disc space/foramina. After the skin and fascia was opened, muscle layers were explored with the help of special speculum-like retractors. After confirming a disc space, the upper part of the lower lamina was drilled first beginning from the upper part of the lower lamina using a high-speed drill under the operating microscope. Thus, a keyhole shaped foraminotomy was carried out, 1/3 in the upper lamina and 2/3 in the lower lamina (Figure 1A). The ligamentum flavum was excised *en masse* with the epidural venous plexus by coagulating the former with a bipolar cautery in order to minimize bleeding from the epidural venous plexus. Moreover, the dural sac and the nerve root were exposed, and the root was gently retracted from its axillary part to reach the extruded disc and to remove the sequestered disc fragment. After confirming that the nerve root and neural structures were decompressed at the foramina and disc space, hemostasis was achieved, and that marked the end of the surgical procedure (Video 1).

PC-EAD; The surgical procedure was begun after the patient had been prepared as in the PC-MD technique. We used Easy-GO® system (Karl-Storz, Tuttingen, Germany) for PC-EAD procedure. Following skin incision, the layers were explored by dilating them with the help of tube-shaped dilators. After confirming the target disc level under fluoroscopy, the procedure was continued under endoscopic view with a 15-mm diameter working canula. In the PC-EAD technique, bone removal from the upper lamina was not needed, thanks to the change of direction of the working cannula and its angled vision. Thus, the upper 2/3 of the lower lamina was first drilled, followed by the upper lamina as deemed necessary by a visual assessment of the surgeon (Figure 1B). The ligamentum flavum was coagulated within lifting by bipolar cautery. The maneuver involved the coagulation of venous plexus beneath

the flavum and its removal *en masse* with the epidural venous plexus without bleeding. If they were separately removed, extensive bleeding could have occurred. The dural sac and the nerve roots were exposed, and the root was gently retracted from its axillary part to reach the extruded disc and to remove the sequestered disc fragment. After confirming that the nerve root and neural structures were relieved at the disc space, hemostasis was achieved, marking the end of the surgical procedure (Video 2).

Patients in both groups were mobilized approximately 8 hours after surgery and were discharged on the first postoperative day. They were prescribed the wearing of neck collars for only three days after their discharge.

Follow-up

The patients underwent routine clinical follow-up at 3, 6, and 12 months after surgery, and a radiological control was performed at 12 months. Cervical lordotic alignment and adjacent segment degeneration were checked at the radiological follow-up using a radiograph and an MRI at 12 months. Their final control was done via outpatient clinic examination or phone call. A 10-point visual pain scale score was used to rate neck and arm pain, and the PROLO score to assess the final status (Table I) (39).

Statistical Analysis

The Statistical Package for Social Sciences (SPSS® for Windows) software was used to compare the preoperative and postoperative parameters among the two groups. The data set was tested for normality using the Kolmogorov Smirnov test. Non-parametric tests were used for skewed data. The Wilcoxon test was used to compare preoperative and postoperative values. The Mann Whitney-U test was used to compare the two techniques to parameters such as duration of operation, volume of bleeding, and postoperative values. A p-value of less than 0.05 was considered statistically significant.

RESULTS

Table II displays the demographic features of patients. Table III summarizes patient characteristics, surgical findings, and postoperative course.

The mean operative time was 74.39 (50–80) min for PC-MD and 81.4 (55–90) min for PC-EAD. Both groups had similar postoperative durations of hospital stay. Volume of bleeding

Table I: Parameters of the Prolo Economic and Functional Scoring

Economic (activity) status		Functional (pain) status	
E1	Complete invalid (worse)	F1	Total incapacity (worse)
E2	No gainful occupation (including housework or retirement activities)	F2	Moderate-to-severe daily pain (no change)
E3	Working/active but not at premorbid level	F3	Low level of daily pain (improved)
E4	Working/active at previous level w/limitation	F4	Occasional or episodic pain
E5	Working/active at previous level w/o restrictions	F5	No pain

Table II: Demographic Characteristics of the Patients were Showed at the Table

	PS – MD	PS – EAD
Age	51.1 (30-70)	38.7 (18-58)
Female	22 (52.3%)	26 (63.4%)
Male	20 (47.7%)	15 (37.6%)
Weakness	26 (61.9%)	20 (48.8%)
Numbness	35 (83.3%)	31 (75.6%)
Distribution of Level	C3-C4	1 (2.4%)
	C4-C5	1 (2.4%)
	C5-C6	21 (50%)
	C6-C7	19 (45.2%)
	C7-T1	-

Table III: Operative Findings and Complications were Seen at the Table

	PC – MD	PC – EAD	p
Operation time (minute)	74.39 (50-80)	81.4 (55-90)	p>0.5
Bleeding (ml)	88.29 (80-100)	81.5 (70-90)	p>0.5
Neurologic deficit	1 (2.3%)	-	
Persistent pain	1 (2.3%)	1 (2.4%)	
CSF leakage	3 (7.1%)	3 (7.3%)	
Recurrent	0	1 (2.4%)	

was also similar between the two techniques (PC-MD; 88.29 mL and PC-EAD; 81.5 mL).

Postoperative VAS and PROLO scores showed significant improvement over preoperative values, but outcomes were similar at the both groups. VAS score for neck pain in PC-MD group was slightly more than in PC-EAD cases, though not significantly different between them (Table IV). Total PROLO score were 9 and 10 points in 75 (90.4%) patients, and rate of the best success group was 90.3%, but 3 cases (3.6%) experienced moderate and/or poor scores. No severe permanent postoperative neurological complications occurred. CSF leak developed in three patients in the PC-MD group and three patients in the PC-EAD group; this complication was managed by administering fibrin glue into the operative field perioperatively. They were later discharged without any neurological sign or wound site complication, and they suffered no complications at follow-up. One patient from each group had persistent postoperative pain, and a control MRI examination showed complete disc removal and full decompression of the nerve root. Such patients received physical therapy followed by gabapentin, which improved their symptoms. One patient in the PC-EAD group had recurrence of pain five months after surgery. An MRI scan revealed a recurrent CDH that was treated by ACDF.

Among patients who were initially treated with PC-EAD, three switched to PC-MD and were analyzed in the PC-MD group. In two of them, the change was made due to an inadequate view of the surgical field owing to an epidural venous bleeding. The other patient was switched to PC-MD because of the entry of a dilator into the spinal canal through the interlaminar space while placing the dilator tubes. That patient developed left-sided postoperative hemiparesis, but regained

Table IV: Comparison of Preoperative and Postoperative Rate of Outcome Scoring (A: PROLO score, B: VAS score) Data Between PS-MD and PS-EAD was Shown at the Tables

All patients		MD		EAD	
PROLO _{preop}	PROLO _{postop}	PROLO _{preop}	PROLO _{postop}	PROLO _{preop}	PROLO _{postop}
7.44	9.62	7.41	9.58	7.44	9.66
p≤0.05		p≤0.05		p≤0.05	
p>0.980					

All patients				MD				EAD			
VAS _{preop} NECK	VAS _{postop} NECK	VAS _{preop} ARM	VAS _{postop} ARM	VAS _{preop} NECK	VAS _{postop} NECK	VAS _{preop} ARM	VAS _{postop} ARM	VAS _{preop} NECK	VAS _{postop} NECK	VAS _{preop} ARM	VAS _{postop} ARM
7.74	2.23	8.78	1.07	7.72	2.32	8.83	1.11	7.75	2.18	8.72	0.97
p≤0.05		p≤0.05		p≤0.05		p≤0.05		p≤0.05		p≤0.05	
p>0.412											

his preoperative motor strength after being treated with postoperative physiotherapy. Although numbness persisted thereafter, he returned to his previous work after treatment with pregabalin.

DISCUSSION

PCD via posterior keyhole foraminotomy could be performed with microscopic or endoscopic systems and is effective in about 93.6% of patients (6,11,19,35). Current surgical tools minimize muscle injury, ensure a lower rate and severity of postoperative pain, and shorten the time of discharge and full recovery. Fusion is unnecessary, so implant complications and pseudarthrosis risk are both eliminated, and permits a natural-like aging process of spine with less risk of adjacent segmental disease (1,10).

Anterior Versus Posterior Approaches

ACDF is associated with several complications including neurological deterioration, dysphagia, esophageal perforation, hoarseness, vascular injury, Horner syndrome and postoperative hematoma (44). Vascular injuries may involve embolic stroke episodes originating from ruptured atheroma plaques due to excessive manipulation of the carotid arteries (16). Esophageal perforation is rare (0.25%) but is associated with high mortality and morbidity due to mediastinitis, abscess, and tracheo-esophageal fistula formation (17). The leading long-term complications of anterior cervical discectomy are related to implants and they include collapse, displacement or failure of fusion (29,32). Dislocation of a cage or bone graft has an incidence of 8%, failure of fusion in 5% for single-level, and 15% for multilevel discectomies (27,29,32,37,44). At 10-year follow-up, 25% of patients were admitted to the hospital with adjacent segmental disease, of whom 2.9% each year required reoperation (18). Moreover, Goffin et al. reported that radiological adjacent segmental disease with or without symptoms developed in 92% of patients after 5 years (15). Biome-

chanical alterations are likely to be responsible for adjacent segmental disease and motion-sparing treatment modalities have been proposed for prevention (8,41). Although biomechanical studies have shown that the use of disc prostheses preserves motion (46), heterotopic ossification (9.4%–17.8%) and spontaneous fusion (2.9%–11%) during follow-up (23,26). ACDF is a particularly risky technique for certain professions such as vocalists, soldiers and athletes (19).

Anterior and posterior approaches can achieve similar success rates with microdiscectomy and endoscopic approaches (6,11,19,35). In a study comparing anterior and posterior endoscopic surgeries, no significant difference was observed between the two groups, although posterior endoscopic surgery was more effective for treatment of sequestered discs (47). No significant difference was reported between ACDF and keyhole foraminotomy in terms of clinical effectiveness and complications (14). A meta-analysis revealed that both techniques had similar clinical effectiveness, while complications were relatively more common in the ACDF group (ACDF - 7% / PCD 4%) and reoperation rates were more common in the PCD group (ACDF - 4% / PCD 6%), although these differences were not significant. Moreover, patients in the PCD group experienced more adjacent segmental disease and less implant procedure, which was an overall advantage (24).

Case Selection for PCD

Laterally sequestered/migrated discs are more suitable for PCD, particularly its endoscopic variety (2). Wen et al. reported that an extruded/sequestered mass at an average distance of 5.41 (1.40) mm from the dural sac was safe and ensured successful outcomes (43). Medially located CDH must be operated via anterior approach, though posterior techniques are indicated for selected cases (14).

The sequester is easily removed if it migrated inferiorly through axilla. Superiorly migrated disc could also be removed by the




Figure 2: Keyhole foraminotomy included two parts as follows; 1/3 of in the upper lamina and 2/3 in the lower lamina was removed that showed by red arms (A), lower part of keyhole was generally enough for removing of sequester in PC-EAD technique, bone windows was showed by green arm (B).

nerve-hook maneuver; however, surgeon should extend to upper part of keyhole and check the shoulder of the nerve root if it descended to more than half of the vertebral body (or is not certain if it could be removed completely).

The consistency of a herniated disc is also important as it is not easy to remove calcified hard discs accompanied by osteophytes via the posterior approach. Moreover, the disc level is completely covered by nerve root due to cranial trajectory of the unco-vertebral joint. Placing a surgical tool in a disc space with such a direction would require significant retraction of the nerve root, thereby leading to compression and injury of the latter (Figure 3). Hence, discs with soft formation that can be removed without vigorous manipulation are suitable to undergo PCD. The procedure performed in this approach entails the removal of a sequestered/migrated part (sequestrectomy). Since compression is anteriorly directed in hard discs accompanied by osteophytes, the posterior approach is an indirect decompression. On the other hand, the height of the intervertebral foramen is also reduced as a result of collapsing disc space during the degenerative process. In our opinion, removal of osteophytes by anterior discectomy allows for direct decompression of cervical disc hernias accompanied by osteophytes while the disc space is elevated by placing an interbody cage, which further enlarges the foramen.

In our clinic, ACDF was preferred for CDH cases with degenerative changes (osteophytes, foraminal stenosis, and multilevel cervical discectomies). PCD is not performed for calcified/hard discs. Similarly, Kim et al. performed an endoscopy-assisted technique for foraminal soft discs and excluded patients with spinal cord compression, and facet joint degeneration or a calcified disc (20). Reutten et al. considered the presence of a foraminal or lateral disc as an indication for surgery among 87 patients operated with PCD via full endoscopic approach; they excluded patients with instability and deformity, a medially located disc, or PLL calcification (33). At a follow-up duration of 24 months, they reported 96.6% clinical success. However, our series had a clinical success rate of 90.3% at a 24.7 months follow-up.

Biomechanics

PCD is a motion-sparing technique that employs minimally-invasive approaches (1,10). Studies on degenerative processes and vertebral alignment after PCD evaluated disc heights and sagittal alignments through control images taken during follow-up visits. No significant reduction in disc height or progression to kyphosis at the sagittal alignment was observed, and segmental motion was maintained. The authors also admitted the shortness of the follow-up time, emphasizing the need for studies with longer follow-up time to be undertaken (20,21).

A biomechanical comparison of anterior cervical discectomy and keyhole foraminotomy suggests that neither of them causes overt instability; in anterior discectomy, however, an approximately two-fold increase occurs in segmented motion and a need for stabilization may arise later (9). Yuchi et al., in a biomechanical study based on the finite element method, reported that segmental motion significantly increased after anterior endoscopic foraminotomy compared with both the control and posterior keyhole foraminotomy groups. Although a slight increase occurred in the posterior keyhole foraminotomy group compared to the control group, it was not significant (49). Generally, anterior approaches cause enhanced FSU motion, which requires fusion. Hence, anterior approaches inevitably progress to ACDF. It has been reported that keyhole foraminotomy and anterior approaches provide similar degrees of foraminal dilation, and no increase occurs in segmental range of motion among patients undergoing keyhole foraminotomy (28). Similarly, an analysis of our patients' preoperative and postoperative MRI findings at a mean follow-up time of 24.7 months did not reveal any degenerative change on the existing findings at the adjacent disc spaces.

Adjacent segmental degeneration is a long-term complication of ACDF, whereas PCD procures preservation of motion as fusion and implants are not used. Alvin et al. reported that ACDF and the posterior keyhole approach were similar in terms of clinical efficacy but ACDF was economically burdensome (3). Another study revealed an increased cost associated with the use of implants (25).




Figure 3: Illustrative drawings showed that relationship nerve root, disc, and disc space is sensitive and rigor, because uncovertebral joint turned to upwards at the foramen and covered by nerve root; **A)** demonstrated the enveloping of them via oblique and laterally perspectives, **B)** illustrated a herniated disc through posterior approach and the extruded fragment must be removed by a nerve hook via pulling it from the underneath without impolite manipulation.

Complications

A learning curve is an inevitable aspect in endoscopic approaches (2,5,40). We experienced a case of dural injury when a dilator tube entered into the canal through the interlaminar space while placing dilator tubes. We then switched to microdiscectomy in order to get a clearer view of the nerve root and spinal cord, and to check the need for repair. That patient suffered a temporary postoperative hemiparesis and persistent sensory complaints; he was able to return to work following gabapentin treatment. In two patients, the operation initially began with PC-EAD and switched later to PC-MD because of a poor view of the surgical field owing to venous bleeding. The distinction between the epidural venous plexus and ligamentum flavum is usually difficult, and venous bleeding is often encountered during flavectomy in cervical surgery. Positioning a patient in the prone position obstructs venous return and complicates stoppage of such bleeding. Therefore, the flavum was coagulated with bipolar cautery and removed in both PC-MD and PC-EAD approaches. However, a switch was made to the PC-MD technique in two of our patients at the beginning of our series.

Another complication seen in our series was a CSF leak. CSF leaks typically occur as a result of dural injury at an axilla of a nerve root. PCD allows the removal of the disc from the injured nerve root. In the axilla, dural repair with primary suturing is challenging due to its direction and anatomical location. The risk of injury to nerve fibers passing by that location during suturing could be a possible threat. Some authors have reported that the dura can be repaired with tools such as the u-clip used in abdominal laparoscopic procedures (30). However, it is troublesome to seal the injury usually developing in the antero-inferior direction at the axillary part of a nerve root with a favorable angle. Likewise, patches of muscular and/or fatty tissue are utilized for dural injury repair. It appears worrisome that a graft placed at an axilla of a nerve root causes irritation of the relatively narrow cervical intervertebral root by creating a mass effect. Six patients in our series suffered dural injury; three (7.1%) patients in the PC-MD group and 3 (7.3%) patients in the PC-EAD group developed intraoperative CSF leak that was managed by spraying a fibrin glue into the operative field. Those patients were discharged free of any neurological sign or wound problem and remained well throughout follow-up.

Although recurrent herniation is the major criticism against PCD approaches, several authors reveal that they are not significantly different from ACDF (42). Selvanathan et al., in a comparative study between ACDF and posterior keyhole foraminotomy, two patients from the ACDF group underwent reoperation due to adjacent segmental disease and one patient due to the need for foraminal decompression (2% in all), while 1 (2%) patient from the keyhole foraminotomy group required surgical intervention due to recurrence of symptoms at a 2-year follow-up visit (36). Wu et al., reported a meta-analysis that included 24 studies with posterior endoscopic procedures. They noted a reoperation rate of 4.8%–5.3% (45). Cervical spinal alignment was analyzed for recurrence in patients who were re-operated on posterior discectomy.

Although Kim et al. reported that cervical curvature does not worsen after PCD (21), Zeng et al. presented a case series that included three patients with recurrence or development of progressive kyphosis who later underwent ACDF (50). Initial marked degeneration, bony spur and/or osteophytes were associated with other levels, while cervical kyphosis could cause inadequate decompression and required reoperation (31). The authors suggested ACDF procedure in patients with recurrence after posterior discectomy (31,50). Only one (1.2%) patient in our series was diagnosed of recurrent herniation and was re-operated through ACDF.

Comparison of Posterior Approaches

Marked differences exist between anterior and posterior approaches with respect to patient selection and technical aspects. However, different posterior approaches are fundamentally similar. Thus, their success rates are similar to one another as well; in a meta-analysis, the clinical success rates of posterior approaches was 93.6% for patients who underwent PC-MD and 89.9% PC-EAD, and both techniques were statistically similar. In addition, both techniques appeared similar in terms of complication rates and the need for reoperation (45).

Fessler and Khoo compared 25 endoscopic cases and 25 microscopic keyhole foraminotomy cases; they reported complete recovery rates of 92% and 87% for radicular pain and neck pain, respectively, in the endoscopic group, and 88% and 89%, respectively, in the microscopic group. Both techniques were similar (13). Kim and Kim randomized 19 patients to the microscopic group and 22 to the endoscopy-assisted keyhole foraminotomy group. Though a marked reduction in the length of hospital stay, postoperative analgesic use, and intraoperative bleeding volume were observed, the clinical outcomes were similar in both groups (22). The same study also reported that endoscope-assisted PCD had a complication rate of 0%–4.3% while the corresponding figure for full endoscopic technique was 3%–5%, with both figures being similar (22). PC-MD and PC-EAD have similar success rates via posterior approaches according to the PROLO and VAS scores; however, the VAS for neck pain in the PC-MD group was slightly more than in the PC-EAD group. We hypothesize that this difference was related to the dilatation of muscle through incision of speculum-like retractor.

Cervical sequestrectomy in posterior approaches is usually performed from the axilla of a nerve root. The ability to perform keyhole foraminotomy with both PC-MD and PC-EAD offers a surgeon the advantage of using both hands. Keyhole foraminotomy does not only provide a corridor for sequestrectomy, but also for an adequate decompression at the foramina. By this way, the nerve root can be retracted gently, and the disc can be removed by a simple draw-aside maneuver using micro tools. It is clear that such fine manipulations would be safer than those carried out using working cannula in pure endoscopic systems. Likewise, disc removal is achieved by retracting the nerve root with invasively-working cannula during posterior full endoscopic cervical discectomy. Therefore, numbness is more common after surgical procedures performed with full endoscopic

systems (45); we believe that the aforementioned temporary sensory complaints were related to that retraction maneuver.

Reutten et al. reported that full endoscopic approaches in posterior cervical techniques do not directly aim at the disc as in lumbar pathologies, although laminoforaminotomy is performed to a lesser degree. He stated that the aim of posterior approaches is to avoid surgical trauma under a good view of the surgical field, and added that full endoscopic procedures fall short in the posterior cervical region. He drew attention to the term “full endoscopic” for techniques applied transdiscally, without necessitating bone resection as is the case for lumbar and anterior cervical approaches (33,34). Yet, cervical disc hernias are underneath the nerve root and must be removed from the axillary direction. Thus, keyhole foraminotomy offers an important advantage for safe disc removal via PS-MD and PS-EAD. We therefore postulate that the similar clinical success rates encountered in both techniques could be linked to the fact that keyhole foraminotomy was feasible in both of them.

Limitations

The first limitation of our study includes the use of a single institution and surgeon specialized in microneurosurgery and endoscopic procedures. This could lead to bias for case selection. Secondly, the study was a nonrandomized retrospective one. Finally, the follow-up period was short. Thus, we recommend further studies with prospective, multicentric, and randomized designs with long-term follow-up.

CONCLUSION

A sequestered/migrated disc can be removed using a minimally-invasive technique via keyhole foraminotomy using either PC-MD or PC-EAD approaches with favorable outcomes in selected cases. Minimally-invasive posterior approaches provided keyhole foraminotomy; thus, a surgeon skilful with both hands also enhanced this surgical window. These approaches avert the natural aging process by least affecting muscle tissue, avoiding bone fusion, and/or implants.

AUTHORSHIP CONTRIBUTION

Study conception and design: AD, RK

Data collection: RK, DKG, EI, GG

Analysis and interpretation of results: AD, RK, DKG, ADB

Draft manuscript preparation: AD, RK; DKG, ADB

Critical revision of the article: AD, RK

Other (study supervision, fundings, materials, etc...): AD, RK, DKG

All authors (AD, RK, DKG, EI, GG, ADB) reviewed the results and approved the final version of the manuscript.

REFERENCES

1. Acikbas SC, Ermol C, Akyuz M, Tuncer R: Assessment of adjacent segment degeneration in and between patients treated with anterior or posterior cervical simple discectomy. *Turk Neurosurg* 20:334-340, 2010
2. Ahn Y: Endoscopic spine discectomy: Indications and outcomes. *Int Orthop* 43:909-916, 2019
3. Alvin MD, Lubelski D, Abdullah KG, Whitmore RG, Benzel EC, Mroz TE: Cost-utility analysis of anterior cervical discectomy and fusion with plating (ACDFP) versus posterior cervical foraminotomy (PCF) for patients with single-level cervical radiculopathy at 1-year follow-up. *Clin Spine Surg* 29:E67-E72, 2016
4. Ansari D, Burley HEK, von Glinski A, Elia C, Chapman JR, Oskouian RJ: The new onset of dysphagia four years after anterior cervical discectomy and fusion: Case report and literature review. *Surg Neurol Int* 11:32, 2010
5. Burkhardt BW, Oertel JM: The learning process of endoscopic spinal surgery for degenerative cervical and lumbar disorders using the EasyGO! System. *World Neurosurg* 119:479-487, 2018
6. Bydon M, Mathios D, Macki M, de la Garza-Ramos R, Sciubba DM, Witham TF, Wolinsky JP, Gokaslan ZL, Bydon A: Long-term patient outcomes after posterior cervical foraminotomy: An analysis of 151 cases. *J Neurosurg Spine* 21:727-731, 2014
7. Caglar YS, Bozkurt M, Kahilogullari G, Tuna H, Bakir A, Torun F, Ugur HC: Keyhole approach for posterior cervical discectomy: Experience on 84 patients. *Minim Invas Neurosurg* 50:7-11, 2007
8. Carrier CS, Bono CM, Lebl DR: Evidence-based analysis of adjacent segment degeneration and disease after ACDF: A systematic review. *Spine J* 13:1370-1378, 2013
9. Chen BH, Natarajan RN, An HS, Andersson GB: Comparison of biomechanical response to surgical procedures used for cervical radiculopathy: Posterior keyhole foraminotomy versus anterior foraminotomy and discectomy versus anterior discectomy with fusion. *J Spinal Disord* 14:17-20, 2001
10. Coric D, Adamson T: Minimally invasive cervical microendoscopic laminoforaminotomy. *Neurosurg Focus* 25: E2, 2008
11. Dunn C, Moore J, Sahai N, Issa K, Faloon M, Sinha K, Hwang KS, Emami A: Minimally invasive posterior cervical foraminotomy with tubes to prevent undesired fusion: A long-term follow-up study. *J Neurosurg Spine* 29:358-364, 2018.
12. Epstein JA, Janin Y, Carras R, Lavine LS: A comparative study of the treatment of cervical spondylotic myeloforaminotomy. Experience with 50 cases treated by means of extensive laminectomy, foraminotomy, and excision of osteophytes during the past 10 years. *Acta Neurochir (Wien)* 61:89-104, 1982
13. Fessler RG, Khoo LT: Minimally invasive cervical microendoscopic foraminotomy: An initial clinical experience. *Neurosurgery* 51 Supplement:S37-45, 2002
14. Foster MT, Carleton-Bland NP, Lee MK, Jackson R, Clark SR, Wilby MJ: Comparison of clinical outcomes in anterior cervical discectomy versus foraminotomy for brachialgia. *Br J Neurosurg* 33:3-7, 2019
15. Goffin J, van Loon J, Van Calenbergh FV, Plets C: Long-term results after anterior cervical fusion and osteosynthetic stabilization for fractures and/or dislocations of the cervical

- spine. *J Spinal Disord* 8:500-508, 1995
16. Graffeo CS, Puffer RC, Wijdicks EF, Krauss WE: Delayed cerebral infarct following anterior cervical discectomy and fusion. *Surg Neurol Int* 7:86, 2016
 17. Halani SH, Baum GR, Riley JP, Pradilla G, Refai D, Rodts GE Jr, Ahmad FU: Esophageal perforation after anterior cervical spine surgery: A systematic review of the literature. *J Neurosurg Spine* 25:285-291, 2016
 18. Hilibrand AS, Carlson GD, Palumbo MA, Jones PK, Bohlman HH: Radiculopathy and myelopathy at segments adjacent to the site of a previous anterior cervical arthrodesis. *J Bone Joint Surg (Am)* 81:519-528, 1999
 19. Kim CH, Chung CK, Kim HJ, Jahng TA, Kim DG: Early outcome of posterior cervical endoscopic discectomy: An alternative treatment choice for physically/socially active patients. *J Korean Med Sci* 24:302-306, 2009
 20. Kim CH, Kim KT, Chung CK, Park SB, Yang SH, Kim SM, Sung JK: Minimally invasive cervical foraminotomy and discectomy for laterally located soft disc herniation. *Eur Spine J* 24:3005-3012, 2015
 21. Kim CH, Shin KH, Chung CK, Park SB, Kim JH: Changes in cervical sagittal alignment after single-level posterior percutaneous endoscopic cervical discectomy. *Glob Spine J* 5:31-38, 2015
 22. Kim KT, Kim YB: Comparison between open procedure and tubular retractor assisted procedure for cervical radiculopathy: Results of a randomized controlled study. *J Korean Med Sci* 24:649-653, 2009
 23. Leung C, Casey AT, Goffin J, Kehr P, Liebig K, Lind B, Logroscino C, Pointillart V: Clinical significance of heterotopic ossification in cervical disc replacement: A prospective multicenter clinical trial. *Neurosurgery* 57:759-763; discussion 759, 2005
 24. Liu WJ, Hu L, Chou PH, Wang JW, Kan WS: Comparison of anterior cervical discectomy and fusion versus posterior cervical foraminotomy in the treatment of cervical radiculopathy: A systematic review. *Orthop Surg* 8:425-431, 2016
 25. Mansfield HE, Canar WJ, Gerard CS, O'Toole JE: Single-level anterior cervical discectomy and fusion versus minimally invasive posterior cervical foraminotomy for patients with cervical radiculopathy: A cost analysis. *Neurosurg Focus* 37:E9, 2014
 26. Murrey D, Janssen M, Delamarter R, Goldstein J, Zigler J, Tay B, Darden B: Results of the prospective, randomized, controlled multicenter Food and Drug Administration investigational device exemption study of the ProDisc-C total disc replacement versus anterior discectomy and fusion for the treatment of 1-level symptomatic cervical disc disease. *Spine* 9:275-286, 2009
 27. Nanda A, Sharma M, Sonig A, Ambekar S, Bollam P: Surgical complications of anterior cervical discectomy and fusion for cervical degenerative disk disease: A single surgeon's experience of 1,576 patients. *World Neurosurg* 82:1380-1387, 2014
 28. Nguyen J, Chu B, Kuo CC, Leasure JM, Ames C, Kondrashov D: Changes in foraminal area with anterior decompression versus keyhole foraminotomy in the cervical spine: A biomechanical investigation. *J Neurosurg Spine* 27:620-626, 2017
 29. Noordhoek I, Koning MT, Jacobs WCH, Vleggeert-Lankamp CLA: Incidence and clinical relevance of cage subsidence in anterior cervical discectomy and fusion: A systematic review. *Acta Neurochir (Wien)* 160:873-880, 2018
 30. Oertel JM, Burkhardt BW: Full endoscopic treatment of dural tears in lumbar spine surgery. *Eur Spine J* 26:2496-2503, 2017.
 31. Oh HS, Hwang BW, Park SJ, Hsieh CS, Lee SH: Percutaneous endoscopic cervical discectomy (PECD): An analysis of outcome, causes of reoperation. *World Neurosurg* 102:583-592, 2017
 32. Pirkle S, Kaskovich S, Cook DJ, Ho A, Shi LL, Lee MJ: Cages in ACDF are associated with a higher nonunion rate than allograft: A stratified comparative analysis of 6130 patients. *Spine (Phila Pa 1976)* 44:384-388, 2019
 33. Ruetten S, Komp M, Merk H, Godolias G: A new full-endoscopic technique for cervical posterior foraminotomy in the treatment of lateral disc herniations using 6.9-mm endoscopes: Prospective 2-year results of 87 patients. *Minim Invas Neurosurg* 50:219-226, 2007
 34. Ruetten S, Komp M, Merk H, Godolias G: Full-endoscopic cervical posterior foraminotomy for the operation of lateral disc herniations using 5.9 mm endoscopes. A prospective, randomized, controlled study. *Spine (Phila Pa 1976)* 33:940-948, 2008
 35. Sahai N, Changoor S, Dunn CJ, Sinha K, Hwang KS, Faloon M, Emami A: Minimally invasive posterior cervical foraminotomy as an alternative to anterior cervical discectomy and fusion for unilateral cervical radiculopathy: A systematic review and meta-analysis. *Spine (Phila Pa 1976)* 44:1731-1739, 2019
 36. Selvanathan SK, Beagrie C, Thomson S, Corns R, Deniz K, Derham C, Towns G, Timothy J, Pal D: Anterior cervical discectomy and fusion versus posterior cervical foraminotomy in the treatment of brachialgia: The Leeds spinal unit experience (2008–2013). *Acta Neurochir (Wien)* 157:1595-1600, 2015
 37. Tasiou A, Giannis T, Brotis AG, Siasios I, Georgiadis I, Gatos H, Tsianaka E, Vagkopoulos K, Paterakis K, Fountas KN: Anterior cervical spine surgery-associated complications in a retrospective case-control study. *J Spine Surg* 3:444-459, 2017
 38. van Limbeek J, Jacobs WC, Anderson PG, Pavlov PW: A systematic literature review to identify the best method for a single level anterior cervical interbody fusion. *Eur Spine J* 9:129-136, 2000
 39. Vanti C, Prosperi D, Boschi M: The prolo scale: History, evolution and psychometric properties. *J Orthop Traumatol* 14:235-245, 2013
 40. Wang B, Lü G, Patel AA, Ren P, Cheng I: An evaluation of the learning curve for a complex surgical technique: The full endoscopic interlaminar approach for lumbar disc herniations. *Spine J* 11:122-130, 2011
 41. Wang F, Hou HT, Wang P, Zhang JT, Shen Y: Symptomatic adjacent segment disease after single-level anterior cervical discectomy and fusion: Incidence and risk factors. *Med*

- (Baltim) 96: e8663, 2017
42. Wang TY, Lubelski D, Abdullah KG, Steinmetz MP, Benzel EC, Mroz TE: Rates of anterior cervical discectomy and fusion after initial posterior cervical foraminotomy. *Spine J* 15:971-976, 2015
 43. Wen H, Wang X, Liao W, Kong W, Qin J, Chen X, Lv H, Friis T: Effective range of percutaneous posterior full-endoscopic paramedian cervical disc herniation discectomy and indications for patient selection. *BioMed Res Int* 2017: 3610385, 2017
 44. Wilson JR, Radcliff K, Schroeder G, Booth M, Lucasti C, Fehlings M, Ahmad N, Vaccaro A, Arnold P, Sciubba D, Ching A, Smith J, Shaffrey C, Singh K, Darden B, Daffner S, Cheng I, Ghogawala Z, Ludwig S, Buchowski J, Brodke D, Wang J, Lehman RA, Hilibrand A, Yoon T, Grauer J, Dailey A, Steinmetz M, Harrop JS: Frequency and acceptability of adverse events After anterior cervical discectomy and fusion: A survey study From the Cervical Spine Research Society. *Clin Spine Surg* 31: E270-277, 2018
 45. Wu PF, Li YW, Wang B, Jiang B, Tu ZM, Lv GH: Posterior cervical foraminotomy via full-endoscopic versus microendoscopic approach for radiculopathy: A systematic review and meta-analysis. *Pain Phys* 22:41-52, 2019
 46. Xu S, Liang Y, Zhu Z, Qian Y, Liu H: Adjacent segment degeneration or disease after cervical total disc replacement: A meta-analysis of randomized controlled trials. *J Orthop Surg Res* 13:244, 2018
 47. Yang JS, Chu L, Chen L, Chen F, Ke ZY, Deng ZL: Anterior or posterior approach of full-endoscopic cervical discectomy for cervical intervertebral disc herniation? A comparative cohort study. *Spine (Phila Pa 1976)* 39:1743-1750, 2014
 48. Yee TJ, Swong K, Park P: Complications of anterior cervical spine surgery: A systematic review of the literature. *J Spine Surg* 6:302-322, 2020.
 49. Yuchi CX, Sun G, Chen C, Liu G, Zhao D, Yang H, Xu B, Deng S, Ma X, Du CF, Yang Q: Comparison of the biomechanical changes after percutaneous full-endoscopic anterior cervical discectomy versus posterior cervical foraminotomy at C5-C6: A finite element-based study. *World Neurosurg* 128:e905-911, 2019
 50. Zheng C, Huang X, Yu J, Ye X: Posterior percutaneous endoscopic cervical discectomy: A single-center experience of 252 cases. *World Neurosurg* 120:e63-e67, 2001