



## Original Investigation

# Prevalence of Anomally Originating Occipital Artery in a Group of Turkish Individuals: A Retrospective Study using Angiography

Nese KESER<sup>1</sup>, Osman KULA<sup>2</sup>, Merih IS<sup>1</sup>, Ismail YARDIMCIOGLU<sup>2</sup><sup>1</sup>University of Health Sciences, Istanbul Fatih Sultan Mehmet Education and Research Hospital, Neurosurgery Clinic, Istanbul, Turkey<sup>2</sup>University of Health Sciences, Istanbul Fatih Sultan Mehmet Education and Research Hospital, Radiology Clinic, Istanbul, Turkey

Corresponding author: Nese KESER ✉ nskeser@gmail.com

## ABSTRACT

**AIM:** To investigate the prevalence of anomalously originating occipital artery (OA) using angiography in a group of Turkish individuals.

**MATERIAL and METHODS:** The images recorded in the picture archiving and communication system for a total of 114 patients, in which the whole vertebral artery, as well as the external carotid artery (ECA) and its branches, were visualized, were retrospectively reviewed. Images were obtained using a Toshiba INFNX-i 8000V (Canon Medical Systems, Otawara, Tochigi, Japan) angiography device.

**RESULTS:** We diagnosed 11 cases (12 arteries) with anomalously originating OA, representing a prevalence of 9.64%. In 7 cases, the ascending pharyngeal artery and OA originated with a common root from the ECA (8 arteries); and in 4 patients, OA originated from the distal part of the ECA (C1 vertebral level).

**CONCLUSION:** As for many vascular structures, the prevalence of OA variations may vary according to the population under question and the examination method used. Our study has shown that in a sample from the Turkish population, the prevalence of anomalously originating OA was much higher than that stated in the literature, which used magnetic resonance angiography. To avoid complications, the high prevalence of this anomaly must be taken into account during surgeries that require the use of OA, endarterectomies, and endovascular interventions.

**KEYWORDS:** Anatomy, Angiography, Arterial variations, External carotid artery, Occipital artery

## INTRODUCTION

The occipital artery (OA) usually originates from the posterior aspect of the proximal external carotid artery (ECA) (42,45), and can be divided into three distinct segments: digastric, horizontal (suboccipital), and terminal (occipital) (Figure 1) (3). Based on anatomical, surgical, and radiological studies, variations of OA have been described in the literature (1,3,6,7,9-11,14,15,20-23,27-30,37,38,41-43). It has also been reported in several publications that the

incidence of some cerebrovascular diseases, Willis polygon, and superior thyroid artery variations differ among populations (12,13,17-19,25,31,32,40,44,46).

To the best of our knowledge, no studies, published in English, have investigated the prevalence of OA origin variations using angiography. We investigated and reported the rate and types of OA origin variations using catheter angiography in a group of Turkish individuals.

## ■ MATERIAL and METHODS

All the images recorded in the picture archiving and communication system of the 440 cerebral angiography procedures performed in the Interventional Neuroradiology Unit of Fatih Sultan Mehmet Education and Research Hospital Radiology Department between July 2016 and December 2017 were retrospectively reviewed. A total of 114 patients had the entire vertebral and external carotid arteries, and their branches, visualized. Most of the examined patients had suspected or proven ischemic cerebrovascular disease. Of the 114 patients, 79 were men and 35 were women, and the mean age was 64.3 years (range 41-86 years). The need for informed consent was waived by the internal institutional review board because of the retrospective nature of the study.

Toshiba INFNX-i 8000V (Canon Medical Systems, Otawara, Tochigi, Japan) was used as the angiography device. A total of 12 cc of non-ionic contrast medium (OMNIPAQUE 300 mg iodine/ml, GE Healthcare, Cork, Ireland) was injected at a rate of 4 cc/sec into the corresponding common carotid artery (CCA) to visualize the ECA, OA, and their branches. A total of 5 cc of contrast medium, injected at a rate of 3 cc/sec was used to visualize VA origin and its proximal part, and a total of 9 cc injected through the proximal part of VA at a rate of 3 cc/sec was used to visualize the distal part of vertebral artery (VA) and the posterior circulation. Two experienced neuroradiologists, Murat Velioglu and Osman Kula, reviewed all angiographic images to observe the abnormal origin of OA.

## ■ RESULTS

Eleven cases with anomalously originating OA were detected among the 114 cases (9.64%) using catheter angiography. The ascending pharyngeal artery (APA) and OA originated from the ECA with a common root in 7 of the 11 cases with OA variation. In two of them, the pharyngo-occipital trunk (POT) originated from a proximal part of the ECA (Figure 2), and one had bilateral POT (Figure 3A, B). In 4 patients, OA originated from the distal part of the ECA (C1 vertebral level) and was directed posteriorly (Figure 4).

## ■ DISCUSSION


We found anomalously originating OA in 11 out of 114 cases (9.64%), which was well above the previously reported incidence in studies of OA variation using non-contrast magnetic resonance angiography (MRA) (42). To the best of our knowledge, this is the first report about the incidence of anomalously originating OA using catheter angiography.

Many variations of OA were described in postmortem dissections before 1841 (14,22). Recent reports describing OA origin variations are limited to case reports of incidental angiographic or intraoperative findings (1-3,6,10,11,15,20-22,27-30,33,38,41,43,45). We found only one incidence study using non-contrast MRA in the literature published in English, in which the incidence of OA origin variation was reported as 0.21% (42).

Currently, catheter-based digital subtraction angiography (DSA) is considered the gold standard for the diagnosis of occlusive extra- and intracranial carotid artery diseases (39). Slow, turbulent, and non-turbulent flow cannot be detected in unenhanced MRA (39), so anomalously originating blood vessels may be missed during MRA studies. OAs originating from CCA have also been reported in the literature (15,45), and the true prevalence of this variant may have been underestimated with MRA. Another reason for the lower rate of variations in the study by Uchino et al. is that MRA was performed up to the carotid bifurcation only (42).


CCA usually divides into the ECA and internal carotid artery (ICA) at the C3-4 level (16). The OA classically originates from the posterior part of the ECA at 20 mm distal to the carotid bifurcation, then turns medial to the posterior bundle of the digastric muscle, and after passing through the medial part of the mastoid advances posteriorly between the occipital bone and C1 (Figure 1) (26,45). The exact origin of the OA is over the origin of the facial artery in 57% of individuals (Figure 1), between the facial and lingual arteries in 32%, and below the lingual artery in 11% (16). In one of our cases, OA originated at the C1 level (Figure 4), and we could not find similar reports in our search of the literature published in English.

The embryology of OA and other variant origins of ECA branches were not fully understood until Lasjaunias et al. hypothesized that OA is formed from the proatlantal intersegmental artery (26). The ascending pharyngeal artery (APA) often shares a common origin with OA, and the



**Figure 1:** Photograph showing the classic origin of the occipital artery from the left external carotid artery and the digastric, suboccipital and proximal occipital segments of the occipital artery. \*: origin of the occipital artery; \*\*: facial artery; **EAM:** external auditory meatus; **ECA:** external carotid artery; **ICA:** internal carotid artery; **VA:** vertebral artery.

phylogenetic and embryological details of this common origin of APA and OA has been previously described (1,26). While Hayashi et al. found the prevalence of APA to be 19% in Japanese cadaveric dissections (16), Cavalcanti et al. reported it at 7.9% in their cadaveric dissections (8). In our study using DSA, we detected 7 cases (6.1%) in whom the OA and APA originated by a common root from ECA. While the two POTs were localized proximal to the left ECA (Figure 2), the bilateral



**Figure 2:** Left lateral angiogram is showing the apparently proximal origin of the pharyngo-occipital trunk in relation to the external carotid artery (**large arrow**), followed by the ascending pharyngeal (**arrow**) and occipital arteries (**interrupted arrow**). Also, internal carotid artery dissection is observed (**arrowhead**).

output of the POT was noted in 1 case (0.87%) (Figure 3A, B). To the best of our knowledge, no other cases with bilateral POT have been reported in the literature published in English.


Anomalies of the carotid vessels are usually asymptomatic, but they are increasingly becoming recognized with the increasing number of surgical procedures and endovascular interventions of the carotid arteries and their branches (20). However, these variations recognized by chance may sometimes be clinically important (11,15,30). In the literature, it was reported that a case was mistakenly diagnosed with carotid stenosis using ultrasonography and was planned for stenting, but after DSA, it was found that OA and APA were both originating from an occluded ICA stump, and the intervention was abandoned (43). Also, OA is a commonly used vessel in the revascularization surgery of the posterior cranial fossa as an intracranial bypass graft (4,5,24,34-36). Taking the OA variations into consideration and evaluating it with DSA before bypass surgery is important for surgical planning. Atherosclerotic stenosis of the extracranial carotid artery typically occurs at the junction of the CCA, ECA, and ICA arteries (11,14). Hence extracranial ICA anomalies affect the plan of surgical endarterectomies. In their study, Hayashi et al. emphasized the importance of awareness of the variations in the origin of vessels during carotid endarterectomy to avoid injuries or problematic back bleeds during atherosclerotic plaque extraction (16).

The limitations of our study include the retrospective nature and the small sample size. Moreover, almost all our cases have been diagnosed with or were suspected to have, atherosclerotic vascular disease (Figure 2).

Uchino et al. suggest that since atheromatous plaques are most frequently observed at the posterior aspect of the proximal ICA, OA arising from the posterior aspect of the ICA may be occluded in some of these patients (42). Therefore,



**Figure 3:** Case with the bilateral pharyngo-occipital trunk. **A)** A lateral angiogram performed from the right main carotid artery better delineates the pharyngo-occipital trunk (**arrow**) originating from the posterior wall of the external carotid artery. The ascending pharyngeal artery (**arrowheads**) and the occipital artery (**interrupted arrow**) is followed along its course. **B)** The pharyngo-occipital trunk (**arrow**) originating from the posterior wall of the external carotid artery in a lateral angiogram performed from the left main carotid artery of the same case. The ascending pharyngeal (**arrowheads**) and the occipital arteries (**interrupted arrow**) are followed along with its course.



**Figure 4:** A DSA image from the lateral projection of left common carotid artery injection. Note the occipital artery (**arrow**) is arising from the external carotid artery at C1 vertebral level.

occlusion of the orifice of the OAs may have caused some of our cases to be overlooked, and the rate of variations may have been less than the rate found in anatomic studies. While some of the OA variations are often detected on angiographic images (20-22,30), others require gross anatomic dissections to reveal the relationship between OA and adjacent soft tissues (1,3,10,27,41). However, angiography is more easily performed and is more widespread than anatomical dissections in almost all populations and can be used to evaluate most of the cases.

## CONCLUSION

In our study of Turkish individuals using angiography, we found that the prevalence of anomalously originating OA was much higher than that reported in the literature using MRA. Before surgeries that require the use of OA, endarterectomies and endovascular procedures, a detailed examination is necessary to avoid complications resulting from the high rate of variations.

## ACKNOWLEDGEMENT

We would like to thank Dr. Murat Velioglu for his radiological evaluations and scientific contributions to this project.

## REFERENCES

- Aggarwal NR, Krishnamoorthy T, Devasia B, Menon G, Chandrasekhar K: Variant origin of superior thyroid artery, occipital artery and ascending pharyngeal artery from a common trunk from the cervical segment of internal carotid artery. *Surg Radiol Anat* 28:650-653, 2006
- Altmann F: Anomalies of the internal carotid artery and its branches; their embryological and comparative anatomical significance; report of a new case of persistent stapedia artery in man. *Laryngoscope* 57:313-339, 1947
- Alvernia JE, Fraser K, Lanzino G: The occipital artery: A microanatomical study. *Neurosurgery* 58 Suppl 1:114-122, 2006
- Ates O, Ahmed AS, Niemann D, Baskaya MK: The occipital artery for posterior circulation bypass: Microsurgical anatomy. *Neurosurg Focus* 24(2): E9, 2008
- Ausman JI, Pearce JE, Vacca DF, Diaz FG, Shrontz CE, Patel S: Tandem bypass: Occipital artery to posterior inferior cerebellar artery side-to-side anastomosis and occipital artery to anterior inferior cerebellar artery end-to-side anastomosis-a case report. *Neurosurgery* 22:919-922, 1988
- Benson MT, Hamer JD: Anomalous origin of the occipital artery from the cervical internal carotid artery. *J Vasc Surg* 8:643-645, 1988
- Bowen JC, Garcia M, Garrard CL, Mankin CJ, Fluke MM: Anomalous branch of the internal carotid artery maintains patency distal to a complete occlusion diagnosed by duplex scan. *J Vasc Surg* 26:164-167, 1997
- Cavalcanti DD, Reis CV, Hanel R, Safavi-Abbasi S, Deshmukh P, Spetzler RF, Preul MC: The ascending pharyngeal artery and its relevance for neurosurgical and endovascular procedures. *Neurosurgery* 65 Suppl 6:114-120, 2009
- Cetin N, Akkan K, Ucar M, Onal B, Ilgit E: Bilateral occipital arteries of internal carotid origin: Report of a case and review of the literature. *Journal of the Belgian Society of Radiology* 99: 69-71, 2015
- Chitra R: Trifurcation of the right common carotid artery. *Indian J Plast Surg* 41:85-88, 2008
- Cohen JE, Leker RR, Moshe Gomori J, Itshayek E: Pharyngo-occipital artery variant arising proximal to occluded internal carotid artery: The risk of an unnecessary endarterectomy. *J Clin Neurosci* 21:529-531, 2014
- De Silva KR, Silva R, Amaratunga D, Gunasekera WS, Jayasekera RW: Types of the cerebral arterial circle (circle of Willis) in a Sri Lankan population. *BMC Neurol* 11:5, 2011
- Flaherty ML, Woo D, Haverbusch M, Sekar P, Khoury J, Sauerbeck L, Moomaw CJ, Schneider A, Kissela B, Kleindorfer D, Broderick JP: Racial variations in location and risk of intracerebral hemorrhage. *Stroke* 36:934-937, 2005
- Graham RB, Sugrue PA, Rahme RJ, Batjer HH, Bendok BR: Pharyngo-occipital artery variant arising from the internal carotid artery impacting surgical technique during carotid endarterectomy. *J Neurointerv Surg* 5:e14, 2013
- Gurbuz J, Cavdar S, Ozdogmus O: Trifurcation of the left common carotid artery: A case report. *Clin Anat* 14:58-61, 2001
- Hayashi N, Hori E, Ohtani Y, Ohtani O, Kuwayama N, Endo S: Surgical anatomy of the cervical carotid artery for carotid endarterectomy. *Neurol Med Chir (Tokyo)* 45: 25-29, 2005
- Henderson RD, Eliasziw M, Fox AJ, Rothwell PM, Barnett HJ: Angiographically defined collateral circulation and risk of stroke in patients with severe carotid artery stenosis. North American Symptomatic Carotid Endarterectomy Trial (NASCET) Group. *Stroke* 31:128-132, 2000

18. Hoksbergen AW, Majoie CB, Hulsmans FJ, Legemate DA: Assessment of the collateral function of the circle of Willis: Three-dimensional time-of-flight MR angiography compared with transcranial color-coded duplex sonography. *AJNR Am J Neuroradiol* 24:456-462, 2003
19. Horikoshi T, Akiyama I, Yamagata Z, Sugita M, Nukui H: Magnetic resonance angiographic evidence of sex-linked variations in the circle of Willis and the occurrence of cerebral aneurysms. *J Neurosurg* 96:697-703, 2002
20. Iwai T, Izumi T, Inoue T, Maegawa J, Fuwa N, Mitsudo K, Tohnai I: Occipital artery arising from the anterior aspect of the internal carotid artery identified by three-dimensional computed tomography angiography. *Iran J Radiol* 9:103-105, 2012
21. Iwai T, Izumi T, Inoue T, Maegawa J, Mitsudo K, Tohnai I: Incidence of the occipital artery arising from the internal carotid artery identified by three-dimensional computed tomographic angiography. *Br J Oral Maxillofac Surg* 50:373-375, 2012
22. Jeleu L, Guirov K: Superficial occipital artery: Report of a rare case and literature review. *Eur J Anat* 18:120-122, 2014
23. Kaneko K, Akita M, Murata E, Imai M, Sowa K: Unilateral anomalous left common carotid artery; a case report. *Ann Anat* 178:477-480, 1996
24. Keser N, Avci E, Soylemez B, Karatas D, Baskaya MK: Occipital artery and its segments in vertebral artery revascularization surgery: A microsurgical anatomic study. *World Neurosurg* 112: e534-e539, 2018
25. Kidwell CS, Rosand J, Norato G, Dixon S, Worrall BB, James ML, Elkind MS, Flaherty ML, Osborne J, Vashkevich A, Langefeld CD, Moomaw CJ, Woo D: Ischemic lesions, blood pressure dysregulation, and poor outcomes in intracerebral hemorrhage. *Neurology* 88: 782-788, 2017
26. Lasjaunias P, Théron J, Moret J: The occipital artery. Anatomy-normal arteriographic aspects-embryological significance. *Neuroradiology* 15:31-37, 1978
27. Marques SR, Itezerote AM, Saviolo MR, de Angelis MA, Prates JC: Anatomical variation of the occipital artery. *Rev Chil Anat* 20:193-196, 2002
28. Matsuda I, Handa J, Handa H, Mizuno H: Bilateral anomalous occipital artery of internal carotid origin: Case report. *Nihon Geka Hokan* 46:57-61, 1977
29. Morimoto T, Nitta K, Kazekawa K, Hashizume K: The anomaly of a non-bifurcating cervical carotid artery. Case report. *J Neurosurg* 72:130-132, 1990
30. Newton TH, Young DA: Anomalous origin of the occipital artery from the internal carotid artery. *Radiology* 90:550-552, 1968
31. Nogueira GJ: Pattern of cerebral aneurysms in Morocco: Review of the concept of their rarity in developing countries: Report of 200 cases. *Neurosurgery* 51:849-850, 2002
32. Ongeti KW, Ogeng'o JA: Variant origin of the superior thyroid artery in a Kenyan population. *Clin Anat* 25:198- 202, 2012
33. Ozgur O, Sindel M, Hizay A, Ozturk S, Aytac G, Sindel T: Occipital artery arising from the internal carotid artery: A case report. *Surg Radiol Anat* 39:219-222, 2017
34. Roski RA, Spetzler RF, Hopkins LN: Occipital artery to posterior inferior cerebellar artery bypass for vertebrobasilar ischemia. *Neurosurgery* 10:44-49, 1982
35. Sekhar LN, Natarajan SK, Ellenbogen RG, Ghodke B: Cerebral revascularization for ischemia, aneurysms, and cranial base tumors. *Neurosurgery* 62 Suppl 3:1373-1408, 2008
36. Sundt TM Jr, Piepgras DG: Occipital to posterior inferior cerebellar artery bypass surgery. *J Neurosurg* 48:916-928, 1978
37. Suzuki S, Nobechi T, Itoh I, Yakura M, Iwashita K: Persistent proatlantal intersegmental artery and occipital artery originating from internal carotid artery. *Neuroradiology* 17:105-109, 1979
38. Teal JS, Rumbaugh CL, Segall HD, Bergeron RT: Anomalous branches of the internal carotid artery. *Radiology* 106: 567-573, 1973
39. Thurnher SA: MRA of the carotid arteries. *Eur Radiol* 15 Suppl 5:11-16, 2005
40. Toni R, Della Casa C, Castorina S, Malaguti A, Mosca S, Roti E, Valenti G: A meta-analysis of superior thyroid artery variations in different human groups and their clinical implications. *Ann Anat* 186:255-262, 2004
41. Tubbs RS, Salter G, Oakes WJ: Continuation of the ascending cervical artery as the occipital artery in man. *Anat Sci Int* 79:43-45, 2004
42. Uchino A, Saito N, Mizukoshi W, Okada Y: Anomalous origin of the occipital artery diagnosed by magnetic resonance angiography. *Neuroradiology* 53:853-857, 2011
43. Ustunsoz B, Gumus B, Koksali A, Koroglu M, Akhan O: Missed total occlusion due to the occipital artery arising from the internal carotid artery. *Cardiovasc Intervent Radiol* 30:116-117, 2007
44. White H, Boden-Albala B, Wang C, Elkind MS, Rundek T, Wright CB, Sacco RL: Ischemic stroke subtype incidence among whites, blacks, and Hispanics: The Northern Manhattan Study. *Circulation* 111:1327-1331, 2005
45. Williams PL, Warwick R, Dyson M, Bannister LH (eds), Gray's Anatomy. 37th ed. Edinburgh: Churchill Livingstone, 1989: 734-739
46. Woo D, Rosand J, Kidwell C, McCauley JL, Osborne J, Brown MW, West SE, Rademacher EW, Waddy S, Roberts JN, Koch S, Gonzales NR, Sung G, Kittner SJ, Birnbaum L, Frankel M, Testai FD, Hall CE, Elkind MS, Flaherty M, Coull B, Chong JY, Warwick T, Malkoff M, James ML, Ali LK, Worrall BB, Jones F, Watson T, Leonard A, Martinez R, Sacco RI, Langefeld CD: The ethnic/racial variations of intracerebral hemorrhage (ERICH) study protocol. *Stroke* 44(10):e120-125, 2013